

POSTER PRESENTATION

Open Access

An HPV-E6/E7 immunotherapy plus PD-1 checkpoint inhibition results in tumor regression and reduction in PD-L1 expression

Adrian E Rice^{1*}, Yvette Latchman¹, Joseph P Balint¹, John H Lee², Elizabeth S Gabitzsch¹, Frank R Jones¹

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015) National Harbor, MD, USA. 4-8 November 2015

The prevalence of head and neck cancers in the USA is estimated to be about 370,000 and between 25% to 38% of these are human papilloma virus (HPV)-associated head and neck squamous cell carcinomas (HNSCC). We have developed a viral gene delivery platform to immunize against HPV 16 genes E6 and E7 (Ad5 [E1-, E2b-]-E6/E7) for the treatment of HPV-associated HNSCC. We tested the Ad5 [E1-, E2b-]-E6/E7 immunotherapy alone and in combination with programmed death-ligand 1 (PD-1) blockade in a murine HPV⁺ tumor model. As a single agent, Ad5 [E1-, E2b-]-E6/E7 induced HPV-E6/E7 cell-mediated immunity and resulted in the clearance of small tumors and an overall survival benefit in mice with larger established tumors. When immunotherapy was combined with PD-1 immune checkpoint blockade, an increased level of anti-tumor activity against large tumors was observed in addition to an improvement in survival. Tumor microenvironment analysis in Ad5 [E1-, E2b-]-E6/E7 treated mice revealed an increase in CD8⁺ tumor-infiltrating lymphocytes (TILs). In addition, we observed induction of suppressive mechanisms such as programmed death-ligand 1 (PD-L1) expression on tumor cells and an increase in PD-1⁺ TILs. When Ad5 [E1-, E2b-]-E6/E7 immunotherapy was combined with anti-PD-1 antibody, we observed CD8⁺ TILs at the same increased level but found that a smaller fraction of these were PD-1⁺. Furthermore, we observed a reduction in PD-L1 expression on tumor cells, providing a mechanism by which combination therapy favors tumor clearance and a rationale for pairing antigen-specific vaccines with checkpoint inhibitors in future clinical trials.

Authors' details

¹Etubics Corporation, Seattle, WA, USA. ²Sanford Cancer Research Center, University of South Dakota Medical School, Sioux Falls, SD, USA.

Published: 4 November 2015

doi:10.1186/2051-1426-3-S2-P449

Cite this article as: Rice et al.: An HPV-E6/E7 immunotherapy plus PD-1 checkpoint inhibition results in tumor regression and reduction in PD-L1 expression. *Journal for ImmunoTherapy of Cancer* 2015 **3**(Suppl 2):P449.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Etubics Corporation, Seattle, WA, USA

Full list of author information is available at the end of the article



© 2015 Rice et al. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.