

POSTER PRESENTATION



Genetic engineering of T cells with receptors from NY-ESO-1-specific tumor-recognizing CD4+ T cell as a novel approach for adoptive T cell therapy

Junko Matsuzaki^{1*}, Takemasa Tsuji¹, Immanuel Luescher², Hiroshi Shiku³, Junichi Mineno⁴, Sachiko Okamoto⁴, Lloyd Old⁵, Protul Shrikant⁶, Sacha Gnjatic⁷, Kunle Odunsi¹

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015) National Harbor, MD, USA. 4-8 November 2015

Background

Tumor antigen-specific CD4⁺ T cells generally orchestrate and regulate innate and adaptive immune cells to provide immune surveillance against malignancy. However, activation of antigen-specific CD4⁺ T cells is restricted at local tumor sites where antigen-presenting cells are frequently dysfunctional, which can cause rapid exhaustion of anti-tumor immune responses. Herein, we characterize anti-tumor effects of a unique human CD4⁺ helper T cell subset that directly recognizes the cytoplasmic tumor antigen, NY-ESO-1, presented by MHC class II (MHC-II) on cancer cells. In addition, we clone the TCR gene from tumor-recognizing CD4⁺ T cells (TR-CD4) and test the function of TCR gene-engineered cells.

Methods

NY-ESO-1-specfic CD4⁺ or CD8⁺ T cells were obtained from ovarian cancer patients who received NY-ESO-1 vaccine. Full-length TCR α and β chain genes of TR-CD4 were cloned by 5' RACE PCR. TCR gene was transduced into activated T cells by MSCV-based retroviral vector. The effector function was evaluated against cognate peptide-pulsed target cells or NY-ESO-1⁺MHC-II⁺ cancer cell lines by ELISA, intracellular cytokine staining or CTL assay.

Results

TR-CD4, but not conventional NY-ESO-1-specific CD4⁺ T cells, directly recognized cancer cells in MHC-IIdependent and NY-ESO-1-specific manners. Presentation

¹Roswell Park Cancer Institute, Buffalo, NY, USA

of intracellular NY-ESO-1 on MHC-II by cancer cells required non-classical MHC-II antigen presentation mechanisms. Upon direct recognition of cancer cells, TR-CD4 potently induced IFN- γ -dependent growth arrest in cancer cells. In addition, direct recognition of cancer cells triggers TR-CD4 to provide help to NY-ESO-1-specific CD8⁺ T cells by enhancing cytotoxic activity, and improving viability and proliferation. Notably, the TR-CD4 either alone or in combination with NY-ESO-1-specific CD8⁺ T cells significantly inhibited tumor growth *in vivo* in a xenograft model. Finally, retroviral gene-engineering of polyclonally activated T cells with TCR derived from TR-CD4 successfully produced large numbers of functional TR-CD4.

Conclusions

These observations provide mechanistic insights into the role of TR-CD4 in tumor immunity, and suggest that approaches to utilize TR-CD4 will augment anti-tumor immune responses for durable therapeutic efficacy in cancer patients. Large numbers of TR-CD4 that directly recognize cancer cells and enhance CD8⁺ T cell functions can be generated by gene-engineering with TCR from TR-CD4. Antigen-presenting cell-independent provision of CD4-help by TR-CD4 is especially important to enhance durable CD8⁺ T cell anti-tumor functions at the tumor local site. Adoptive T cell therapy using TR-CD4 in combination with CD8⁺ T cells could be a promising strategy for effective eradication of tumors.

Authors' details

¹Roswell Park Cancer Institute, Buffalo, NY, USA. ²Ludwig Center for Cancer Research, Epalinges, Switzerland. ³Mie University Graduate School of Medicine, Tsu, Japan. ⁴TAKARA BIO INC, Otsu, Japan. ⁵Ludwig Institute for



© 2015 Matsuzaki et al. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http:// creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/ zero/1.0/) applies to the data made available in this article, unless otherwise stated.

Full list of author information is available at the end of the article

Cancer Research, New York, NY, USA. $^6{\rm Mayo}$ Clinic, Scottsdale, AZ, USA. $^7{\rm MSSM}$ New York, NY, USA.

Published: 4 November 2015

doi:10.1186/2051-1426-3-S2-P34

Cite this article as: Matsuzaki *et al.*: Genetic engineering of T cells with receptors from NY-ESO-1-specific tumor-recognizing CD4+ T cell as a novel approach for adoptive T cell therapy. *Journal for ImmunoTherapy of Cancer* 2015 **3**(Suppl 2):P34.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

BioMed Central

Submit your manuscript at www.biomedcentral.com/submit