

# Society for Immunotherapy of Cancer (SITC) clinical practice guideline on immunotherapy for the treatment of urothelial cancer

Matthew D Galsky,<sup>1</sup> Arjun V Balar,<sup>2</sup> Peter C Black,<sup>3</sup> Matthew T Campbell,<sup>4</sup> Gail S Dykstra,<sup>5,6</sup> Petros Grivas,<sup>7,8</sup> Shilpa Gupta,<sup>9</sup> Christopher J Hoimes,<sup>10</sup> Lidia P Lopez,<sup>11</sup> Joshua J Meeks,<sup>12,13</sup> Elizabeth R Plimack,<sup>14</sup> Jonathan E Rosenberg,<sup>15,16</sup> Neal Shore,<sup>17</sup> Gary D Steinberg,<sup>18</sup> Ashish M Kamat<sup>19</sup>

**To cite:** Galsky MD, Balar AV, Black PC, *et al.* Society for Immunotherapy of Cancer (SITC) clinical practice guideline on immunotherapy for the treatment of urothelial cancer. *Journal for ImmunoTherapy of Cancer* 2021;**9**:e002552. doi:10.1136/jitc-2021-002552

Accepted 24 May 2021

## ABSTRACT

A number of immunotherapies have been developed and adopted for the treatment of urothelial cancer (encompassing cancers arising from the bladder, urethra, or renal pelvis). For these immunotherapies to positively impact patient outcomes, optimal selection of agents and treatment scheduling, especially in conjunction with existing treatment paradigms, is paramount. Immunotherapies also warrant specific and unique considerations regarding patient management, emphasizing both the prompt identification and treatment of potential toxicities. In order to address these issues, the Society for Immunotherapy of Cancer (SITC) convened a panel of experts in the field of immunotherapy for urothelial cancer. The expert panel developed this clinical practice guideline (CPG) to inform healthcare professionals on important aspects of immunotherapeutic treatment for urothelial cancer, including diagnostic testing, treatment planning, immune-related adverse events (irAEs), and patient quality of life (QOL) considerations. The evidence- and consensus-based recommendations in this CPG are intended to give guidance to cancer care providers treating patients with urothelial cancer.

## BACKGROUND

Urothelial cancer (a term that encompasses cancers of the bladder, urethra, and upper urinary tract) represents a significant public health concern as the sixth most common type of cancer in the US. In the year 2021, an estimated 83,730 new cases of bladder cancer and 4,190 new cases of cancers of the ureter and other urinary organs will be diagnosed in the US, leading to approximately 18,160 deaths.<sup>1</sup> There is a clear and unmet need for additional therapeutic options that may provide effective disease control outcomes without compromising quality of life (QOL) for patients with urothelial cancer.

For several decades, standard of care (SOC) therapies for urothelial cancer

included surgery, chemotherapy, radiotherapy, and intravesical *Bacillus Calmette-Guérin* (BCG), a form of immunotherapy comprising an attenuated bacterial pathogen to promote antitumor immune responses. In recent years, however, the US Food and Drug Administration (FDA) has approved a number of immune checkpoint inhibitors (ICIs) for the treatment of urothelial cancer arising from the bladder or other areas of the urinary tract. The ICIs approved for the treatment of urothelial cancer at the time of manuscript preparation include anti-programmed cell death protein-1 (PD-1) agents (nivolumab and pembrolizumab) and anti-programmed death-ligand 1 (PD-L1) agents (atezolizumab, avelumab, and durvalumab). The disease states for which the FDA has approved ICI therapy include non-muscle-invasive bladder cancer (NMIBC) and locally advanced or metastatic urothelial cancer (mUC) (including bladder, urethra, and upper tract urothelial cancers).<sup>2</sup> Trials are ongoing investigating ICIs in earlier disease stages of NMIBC, mUC, and muscle-invasive bladder cancer (MIBC), as well as in the context of novel combination regimens, such as in combination with anti-cytotoxic T lymphocyte antigen-4 (CTLA-4) agents (ipilimumab), chemotherapies, or targeted agents.

Due to the recent clinical adoption of ICIs for bladder cancer, many uncertainties remain regarding the optimal use of these agents, both as monotherapies and in combination with existing or emerging modalities.<sup>3–6</sup> Of note, although antibody-drug conjugates (ADCs) such as enfortumab vedotin (EV) or vicinimum are derived from key protein constituents of humoral immunity, they were not defined as



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Dr Ashish M Kamat;  
akamat@mdanderson.org

Protected by copyright, including for uses related to text and data mining, AI training, and similar technologies.

J Immunother Cancer: first published as 10.1136/jitc-2021-002552 on 15 July 2021. Downloaded from <http://jitc.bmj.com/> on May 20, 2025 at Department GEZ-LTA ErasmusHogeschool.

immunotherapies for the purposes of this manuscript since their primary mechanism of action includes direct cytotoxicity akin to classical chemotherapies as opposed to immune-mediated anti-tumor effects.<sup>78</sup>

The Society for Immunotherapy of Cancer (SITC) previously convened an expert panel to develop a clinical practice guideline (CPG) in 2017 titled, “The Society for Immunotherapy of Cancer consensus statement on immunotherapy for the treatment of bladder carcinoma.”<sup>9</sup> However, immunotherapeutic options for the treatment of urothelial cancer have expanded substantially since the publication of the 2017 guideline. Therefore, in 2020, SITC convened an expert panel to generate updated and expanded evidence- and consensus-based recommendations for the treatment of urothelial cancer with immunotherapy. The expert panel discussed and made recommendations on topics including diagnostic testing, treatment planning, emerging data on investigational immunotherapies, the management of immune-related adverse events (irAEs), and patient QOL. The recommendations in this manuscript are not intended to replace sound clinical judgment and unique patient-based decisions, but to provide healthcare professionals with current, evidence-based guidance on the use of immunotherapy for the treatment of urothelial cancer. The panel focused solely on drugs approved by the FDA; regulatory status, availability, or common clinical practices may differ in other regions. The full series of SITC CPGs can be found on the SITC website (<https://www.sitcancer.org/guidelines>).

## GUIDELINE DEVELOPMENT METHODS

The Institute of Medicine’s (IOM) Standards for Developing Trustworthy Clinical Practice Guidelines were used as a model to develop the recommendations in this manuscript. IOM standards dictate that guideline development is led by a multidisciplinary expert panel using a transparent process where both funding sources and conflicts of interest are readily reported. This CPG is intended to provide guidance and is not a substitute for the professional judgment of individual treating physicians.

### Conflict of interest management

As outlined by IOM standards, all financial relationships of expert panel members that might result in actual, potential, or perceived conflicts of interest were individually reported. Disclosures were made prior to the onset of manuscript development and updated on an annual basis. In addition, panel members were asked to articulate any actual or potential conflicts at all key decision

points during guideline development, so that participants would understand all possible influences, biases, and/or the diversity of perspectives on the panel. Although some degree of relationships with outside interests are to be expected among experts, panel candidates with significant financial connections that may compromise their ability to fairly weigh evidence (either actual or perceived) were not eligible to participate in guideline development.

Recognizing that guideline panel members are among the leading experts on the subject matter under consideration and guideline recommendations should have the benefit of their expertise, any identified potential conflicts of interests were managed as outlined in SITC’s disclosure and conflict of interest resolution policies. As noted in these policies, panel members disclosing a real or perceived potential conflict of interest may be permitted to participate in consideration and decision-making of a matter related to that conflict, but only if deemed appropriate after discussion and agreement by the expert panel.

The financial support for the development of this guideline was provided solely by SITC. No commercial funding was received.

### Recommendation development

Panel recommendations are based on literature evidence, where possible, and clinical experience, where appropriate.<sup>10</sup> Consensus for the recommendations herein was generated by open communication and scientific debate in small- and whole-group settings, surveying and responses to clinical questionnaires, as well as formal voting in consensus meetings.

For transparency, a draft of this CPG was made publicly available for comment during the development process and prior to publication. All comments were evaluated and considered for inclusion into the final manuscript according to the IOM standard.

### Evidence rating

The evidence- and consensus-based recommendations of the panel were refined throughout the development process in order to obtain the highest possible agreement among the experts, however, the minimum threshold was defined as 75% approval among the voting members. Evidence supporting panel recommendations was graded according to the Oxford Centre for Evidence-Based Medicine (OCEBM) Levels of Evidence Working Group “The Oxford Levels of Evidence 2” (2016 version). A summary of the OCEBM grading scale may be found below (table 1). The level of evidence (LE) for a given recommendation is

**Table 1** Summary of “The Oxford Levels of Evidence 2.” (adapted from OCEBM Levels of Evidence Working Group)

Level 1	Level 2	Level 3	Level 4	Level 5
Systematic review or meta-analysis	Randomized trial or observational study with dramatic effect	Non-randomized, controlled cohort, or follow-up study	Case series, case-control, or historically controlled study	Mechanism-based reasoning

expressed in parentheses following the recommendation (eg, LE: 1). Recommendations without an associated LE were based on expert consensus.

## DIAGNOSTIC TESTS AND BIOMARKERS FOR UROTHELIAL CANCER IMMUNOTHERAPY

Biomarkers to predict response to intravesical BCG therapy have remained elusive, with most candidate biomarkers reported from single institutional series lacking subsequent validation.<sup>11,12</sup> The exception is the UroVysion Bladder Cancer Kit (UroVysion Kit), a fluorescence in situ hybridization (FISH) test. A prospective, multicenter validation study in 150 patients confirmed findings from a previous 200 patient single-center study<sup>13</sup> that the UroVysion Kit can stratify the risk of recurrence in patients with high-risk NMIBC receiving BCG therapy.<sup>14</sup> However, the test's performance characteristics and its variance over time make it unsuitable for guiding individual patient management. Instead, the test could possibly be suitable for use in a clinical trial to help randomize patients with a positive UroVysion Kit result after induction BCG to receive additional BCG versus an experimental treatment.<sup>15</sup>

Predicting response to immune checkpoint blockade in patients with mUC is particularly important since only approximately 20% of patients demonstrate an objective response to therapy.<sup>16</sup> Biomarker discovery in this domain has focused mostly on PD-L1 expression by immunohistochemistry (IHC), tumor mutational burden (TMB) by next-generation sequencing (NGS), and RNA-based signatures.<sup>17–19</sup> Only PD-L1 IHC has demonstrated prognostic value,<sup>20–22</sup> although data are lacking to support the predictive power of PD-L1 expression for clinical benefit with immunotherapy for platinum-refractory disease. Additionally, methodological limitations, including poor concordance among approved assays, have led to some confusion about the definition of “PD-L1 positivity,” especially given multiple different scoring systems that include protein expression on tumor and/or immune cells.<sup>23</sup> Four different PD-L1 IHC assays have been approved by the FDA: the companion diagnostics VENTANA PD-L1 (SP142) for atezolizumab and IHC 22C3 pharmDx for pembrolizumab, and the complementary diagnostics VENTANA PD-L1 (SP263) for durvalumab and IHC 28-8 pharmDx for nivolumab. At present, there is no role for PD-L1 testing for immunotherapy selection for platinum-refractory disease.

In some of the single-arm, early phase trials of ICIs in the platinum-refractory disease population, patients with PD-L1-positive tumors demonstrated higher response rates and longer survival than those with PD-L1-negative tumors, measured by immune cell (IC) PD-L1 expression or by tumor cell (TC) PD-L1 expression (SP142 for atezolizumab, SP263 for durvalumab, IHC 73-10 pharmDx for avelumab, IHC 28-8 for nivolumab).<sup>19,24–27</sup> However, two large phase III, randomized, controlled trials, IMvigor211<sup>28</sup> and KEYNOTE-045,<sup>29</sup> demonstrated

that PD-L1 expression by IHC was not significantly associated with overall survival (OS), progression-free survival (PFS), overall response rate (ORR), or duration of response (DOR) in patients with platinum-refractory mUC treated with ICIs.

The phase II, single-arm KEYNOTE-052 trial of first-line pembrolizumab for cisplatin-ineligible patients with mUC showed an improved ORR in patients with high PD-L1 expression by IHC 22C3 assay (defined as combined positive score (CPS) ≥ 10).<sup>30</sup> However, there was no significant correlation between response rates and PD-L1 expression in the IMvigor210 trial of atezolizumab using the SP142 assay (NCT02108652).<sup>31</sup> When both of these treatments received accelerated approval by the FDA for the first-line treatment of cisplatin-ineligible patients with mUC in 2017, the labels did not restrict treatment to patients based on PD-L1 testing results. However, in June 2018, based on the Data and Safety Monitoring Committee review of the IMvigor130<sup>32</sup> and KEYNOTE-361 trials,<sup>33</sup> which compared checkpoint blockade to either standard carboplatin-based or cisplatin-based chemotherapy, both the FDA and the European Medicines Agency (EMA) issued label changes indicating that cisplatin-ineligible patients should only receive first-line atezolizumab or pembrolizumab if their tumors were PD-L1-positive, as determined by approved companion diagnostic assays (at the time of writing, SP142 for atezolizumab and IHC 22C3 for pembrolizumab). The FDA stipulated that carboplatin-ineligible patients may be eligible for ICI therapy regardless of PD-L1 expression. The data that led to these label changes are outlined in further detail in the **Advanced/metastatic urothelial carcinoma** section.

Importantly, both IMvigor130 and KEYNOTE-361 pooled cisplatin-eligible and -ineligible patients in the primary analysis. Exploratory analysis of IMvigor130 showed evidence for benefit with single-agent atezolizumab for cisplatin-ineligible patients with PD-L1-expressing ICs in 5% of the tumor area by SP142.<sup>34</sup> However, CPS ≥ 10 did not enrich for response to pembrolizumab in the choice-of-carboplatin population in exploratory analysis of KEYNOTE-361.<sup>35</sup> Based on the available data, in the first line, chemotherapy-naïve setting, atezolizumab and pembrolizumab remain treatment options for patients with PD-L1-positive tumors deemed ineligible for cisplatin-based chemotherapy (in the US, based on the specific label) and for patients deemed ineligible for any platinum chemotherapy. It is important to note, however, that the evaluation of risks versus benefits for cisplatin-based chemotherapy is defined somewhat arbitrarily and includes considerations that involve patient comorbidities and the clinical disease state for which a patient is being treated.<sup>36</sup> Notwithstanding clinical trials, harmonized definitions are needed to develop therapies for unmet need populations.

Initial chemotherapy followed by maintenance PD-(L)1 blockade results in significantly improved outcomes, demonstrated in randomized trials described in the **Advanced/metastatic urothelial carcinoma** section, and



has largely supplanted the use of pembrolizumab or atezolizumab for the first-line treatment of mUC.<sup>37,38</sup> The OS benefit with maintenance avelumab was observed regardless of PD-L1 status, suggesting that PD-L1 testing does not currently offer clinical utility after chemotherapy in mUC.<sup>38,39</sup>

Evaluation of candidate predictive biomarkers for benefit with checkpoint blockade in adjuvant disease settings is pending the completion of prospective randomized trials.<sup>40–43</sup> The first of these data were reported in abstract format from the phase III open-label IMvigor010 trial, which tested adjuvant atezolizumab versus observation after radical cystectomy or (nephro)ureterectomy for muscle-invasive urothelial carcinoma.<sup>44</sup> No significant difference was observed in disease-free survival (DFS) in the entire study cohort or in patients with PD-L1-positive (IC2/3) tumors. Improvement in DFS in the CheckMate 274 study with adjuvant nivolumab versus placebo after radical cystectomy or (nephro)ureterectomy for muscle-invasive urothelial carcinoma in both the intent-to-treat (ITT) population and in the subset of patients with PD-L1-positive tumors (determined by the IHC 28-8 assay) has also been reported. Notably, although CheckMate 274 demonstrated clinical benefit with adjuvant nivolumab in the all-comers population, the DFS hazard ratio (HR) was smaller for the group of patients with PD-L1-positive tumors than for the ITT population (0.53 [95% CI 0.34 to 0.84] vs 0.70 [95% CI 0.54 to 0.89], respectively).<sup>45</sup>

Tumors with high microsatellite instability (MSI-H), which is a biomarker of mismatch repair deficiency (dMMR), may benefit from ICIs regardless of primary tumor origin.<sup>46,47</sup> Pembrolizumab received accelerated approval by the FDA for the treatment of patients with unresectable or metastatic, MSI-H (typically measured by PCR assay or NGS) or dMMR (typically measured by IHC or NGS) solid tumors that have progressed following prior treatment and who have no satisfactory alternative treatment options based on data from five pooled trials (KEYNOTE-016, KEYNOTE-164, KEYNOTE-012, KEYNOTE-028, and KEYNOTE-158).<sup>3</sup> dMMR is present in approximately 1% of bladder cancers<sup>48,49</sup> and 7%–14% of upper tract urothelial carcinomas, where it is more commonly seen in the context of Lynch syndrome, which occurs due to germline defects.<sup>49–51</sup> Since bladder cancer and upper tract urothelial carcinoma are also enriched for *FGFR3* alterations and erdafitinib (a pan-fibroblast growth factor receptor (FGFR) inhibitor) represents an alternative therapy for post-platinum urothelial carcinoma, it remains to be determined if both *FGFR3* testing and MSI-H/dMMR testing could be used in conjunction to optimally guide treatment selection and therapy sequence. The optimal sequence of ICI versus FGFR inhibition in patients with urothelial carcinoma with *FGFR3* alterations remains undefined but is being explored in prospective studies, including the THOR trial (NCT03390504).

Exploratory analyses of clinical trials in urothelial cancer have suggested that tumors with high TMB may

respond better to immunotherapy than tumors with low TMB,<sup>18,21,22,40</sup> but data are still lacking on the use of TMB for clinical decision-making in urothelial cancer. However, the FDA has granted accelerated approval to pembrolizumab for the treatment of adult and pediatric patients with unresectable or metastatic solid tumors with high TMB ( $\geq 10$  mutations/megabase (mut/Mb)) that have progressed following prior treatment and who have no satisfactory alternative treatment options. This approval was based on the single-arm, phase II KEYNOTE-158 (NCT02628067) trial, which examined 1,050 patients with various solid tumors, and measured TMB using the FoundationOne CDx assay.<sup>52</sup> As with MSI-H/dMMR testing, TMB status could potentially aid in the selection of one treatment over another in platinum-refractory mUC. However, it is important to note that with the potential approval and use of PD-(L)1 inhibitors in earlier settings, this biomarker is likely to have less impact on treatment selection in the future.

### Panel recommendations

- Currently, the evidence does not support routine use of biomarkers to guide BCG therapy in NMIBC. Cystoscopy (with biopsy/transurethral resection (TUR) of bladder tumor as needed), urine cytology, and periodic upper tract imaging should be used to detect recurrence.
- PD-L1 expression by IHC should be used to guide therapy in patients with mUC who are cisplatin-ineligible but eligible for carboplatin. Patients with PD-L1 negative tumors should receive carboplatin-based combination chemotherapy in this setting, while those with PD-L1 positive tumors can receive either immune checkpoint blockade or carboplatin-based chemotherapy (LE: 2). Clinical trial data otherwise does not currently support the use of PD-L1 expression to select patients with platinum-refractory disease for therapy.
- MSI-H/dMMR testing should be considered in patients with upper tract and bladder urothelial cancer, especially for patients of younger age and/or with relevant personal or family history to rule out Lynch syndrome, which has implications for genetic counseling (LE: 3). The presence of MSI should not change the use of ICIs in advanced urothelial cancer.

### NON-MUSCLE-INVASIVE BLADDER CANCER

The SOC for intermediate- and high-risk NMIBC is BCG induction followed by maintenance. Depending on risk category and BCG availability, however, intravesical chemotherapy may be used for induction and the length of maintenance therapy varies. For BCG-unresponsive high-risk NMIBC, pembrolizumab is approved and may be offered to patients after a balanced discussion of risks and benefits. A treatment algorithm depicting management options for different NMIBC risk categories is shown in [table 2](#).

**Table 2** NMIBC immunotherapy treatment algorithm

NMIBC risk category	Management		
Low-risk	BCG not recommended		
Intermediate-risk (BCG available)	BCG†-induction and 1-year maintenance		
Intermediate-risk (BCG unavailable)	Intravesical chemotherapy	If recurrence occurs	BCG†
High-risk*	BCG† induction and 3 years maintenance	If BCG-unresponsive high-risk CIS NMIBC with or without papillary tumors	Pembrolizumab

Individual rows represent treatment decision options that can be followed from left to right horizontally in adjacent columns.

\*Including NMIBC high-risk cases with CIS or papillary tumors.

†BCG should not be administered to patients with active infection or gross hematuria, but BCG may be administered to patients experiencing asymptomatic bacteriuria. Best supportive measures should be employed to ensure that patients receive a full, adequate course of BCG.

BCG, Bacillus Calmette-Guérin; CIS, carcinoma in situ; NMIBC, non-muscle-invasive bladder cancer.

### First-line NMIBC

BCG is a live, attenuated strain of *Mycobacterium bovis* that is administered intravesically as a therapy for NMIBC. BCG has been an important option for the management of NMIBC for more than four decades. BCG treatment depends on the risk category of NMIBC. Low-risk disease is defined as single, primary, low-grade tumors (G1, Ta).<sup>53</sup> For patients with low-risk NMIBC, the SOC treatment is TUR followed by intravesical chemotherapy, without the use of BCG or other immunotherapies.<sup>54</sup> The International Bladder Cancer Group (IBCG) defines intermediate-risk bladder cancer as multiple and/or recurrent low-grade Ta tumors, with specific factors including number of tumors (> one), size of tumors (>3 cm), timing of recurrence (within 1 year), rate of recurrence (> one per year), and prior treatment.<sup>55</sup> High-risk NMIBC is defined as any T1, high-grade (G3) or CIS disease.<sup>53</sup>

In SWOG-8795 (also identified as INT-0094 or EST-1888), 447 patients with NMIBC were administered BCG or mitomycin C as adjuvant therapy following surgical resection. Among the study population, 377 patients had Ta/T1 NMIBC, and median follow-up was 2.5 years. In patients with Ta/T1 NMIBC, mitomycin C treatment led to a recurrence-free survival (RFS) rate of 43%, while those treated with BCG exhibited an RFS rate of 54% (HR 1.41; 95% CI 1.06 to 1.88;  $p=0.017$ ). It was also noted that local and systemic adverse events (AEs) of grade 1 or 2 were more common in patients treated with BCG ( $p=0.003$ ).<sup>56</sup>

During the Nijmegen study, 437 patients were treated with TICE-BCG, the *Rijksinstituut voor Volksgezondheid en Milieuhygiëne* (RIVM) strain of BCG, or mitomycin C as adjuvant therapy following surgical resection of NMIBC (for papillary lesions,  $n=387$ ) or biopsy (for CIS,  $n=50$ ). Despite the fact that the BCG regimen used was suboptimal (ie, no maintenance therapy was given) and the cohort in the mitomycin C arm received monthly maintenance, at 2-year follow-up, the estimated rate of DFS for patients with papillary lesions was similar across groups, at 65% (95% CI 60% to 70%) for mitomycin C treatment,

54% (95% CI 49% to 59%) for TICE-BCG treatment, and 62% (95% CI 57% to 67%) for RIVM-BCG treatment. The differences between treatment arms were not statistically significant.<sup>57</sup>

An analysis of six pooled phase II clinical trials in which 119 patients with CIS received TICE-BCG as first-line therapy (6 weekly instillations for induction, 12 monthly instillations for maintenance) found an ORR of 75.6%, and complete response (CR) rate of 45.4% at a median follow-up of 47 months. The median PFS was estimated at  $\geq 48$  months.<sup>58</sup> On the basis of the clinical trials discussed above, in August 1998, the FDA approved the use of TICE-BCG for the first-line and adjuvant treatment of CIS, and for the adjuvant treatment of Ta and/or T1 papillary tumors of the bladder following TUR.<sup>59</sup> Several meta-analyses have subsequently concluded that BCG prevents, or at least delays, progression to invasive disease in high-risk or intermediate-risk disease. The largest of these meta-analyses analyzed 24 trials ( $n=4,863$  patients) and showed a 27% reduction in the odds of progression (9.8% vs 13.8%; odds ratio (OR) 0.73;  $p=0.001$ ) in patients treated with maintenance BCG compared with either TUR alone or TUR with chemotherapies other than mitomycin C.<sup>60–62</sup> The meta-analyses illustrated that BCG is only superior to mitomycin C in situations where BCG maintenance is provided.<sup>60 62</sup>

Historically, there have been efforts to administer BCG in conjunction with recombinant interferons. However, a meta-analysis demonstrated that BCG alone was associated with lower risk of recurrence in comparison to BCG with interferon- $\alpha$ -2a (relative risk (RR) 0.57; 95% CI 0.39 to 0.82) and to BCG with interferon- $\alpha$ -2b (RR 0.42; 95% CI 0.30 to 0.59).<sup>63</sup>

The standard BCG dosing regimen contains an induction and a maintenance phase, with induction consisting of BCG instillation once a week for 6 weeks and maintenance consisting of repeat instillations at set time intervals (at 3, 6, 12, 18, 24, 30, and 36 months postinduction). Maintenance instillations occur once a week for 3 weeks at each time interval. This dosing regimen (the ‘6+3

regimen') is supported by the results of EORTC-30962 (NCT00002990), a phase III clinical trial that randomized patients to receive full-dose or one-third-dose BCG with 1 year or 3 years of maintenance instillations. In this trial, no differences in toxicity were identified between one-third-dose and full-dose BCG. Further, for intermediate-risk patients (defined as lower than pT1 and lower than G3 in this study), no significant difference in the 5-year DFS rate was observed between 1 year and 3 years of maintenance. For high-risk patients (at least pT1 or G3 disease), 3 years of maintenance was superior to 1 year of maintenance by percentage of disease-free patients at 5 years for patients receiving full-dose BCG (HR 1.61; 95% CI 1.13 to 2.30;  $p=0.0087$ ).<sup>64</sup>

A complicating factor is that BCG is currently subject to ongoing global shortages, which has impacted the ability of healthcare providers to provide BCG therapy to patients.<sup>65 66</sup> In light of this ongoing shortage, several professional and advocacy societies have modified their guidelines for management of NMIBC and advocated a risk-stratified approach toward BCG administration.<sup>67–70</sup> For this reason, current guidelines recommend a risk-stratified schedule of maintenance instillations: while patients with high-risk disease should receive a full 3-year course of maintenance, patients with intermediate-risk disease may receive shortened courses of 1 year of maintenance therapy.<sup>68 71 72</sup>

BCG treatment is associated with a number of potential AEs, which are cumulative over the course of BCG therapy, including cystitis, dysuria, frequency of urination, and, more rarely, infections or systemic side effects. Vigilance is important, since these symptoms overlap with other common AEs in patients receiving treatment for bladder cancer, including urinary tract infections, sensitivity related to urinary catheterization, or overactive bladder. AEs stemming from BCG therapy may be addressed through temporary withholding of BCG, conventional treatments (including systemic steroids,<sup>73 74</sup> non-steroidal anti-inflammatory drugs (NSAIDs),<sup>74</sup> anti-tuberculosis antibiotics,<sup>75</sup> and quinolone antibiotics<sup>76 77</sup>), or permanent withdrawal of BCG for severe toxicity.<sup>78</sup> A proinflammatory response to BCG, leukocyturia, is associated with both an increase in self-reported AEs<sup>79 80</sup> and response to BCG.<sup>81</sup> It may be the case that the occurrence of systemic side effects is indicative of an immune response to BCG and thus could be used as a marker of response to therapy, although more data is required on this subject.

One well-studied method to reduce BCG-related toxicity is dose reduction. Clinical trials have demonstrated that reduced-dose BCG may be efficacious, with significantly less toxicity than full-dose BCG. Long-term follow-up of 499 patients in one prospective trial found that while patients with multifocal tumors derived more benefit from the standard dose than the reduced dose ( $p=0.0151$ ), the cause-specific survival at 5 years did not differ between the two arms ( $p=0.76$ ). Patients who received reduced doses of BCG were significantly

less likely to experience grade  $\geq 3$  toxicity ( $p<0.001$ ).<sup>82</sup> However, in the EORTC-30962 trial, there were no differences in toxicity between the one-third-dose and full-dose arms.<sup>64</sup> Reduced dosing frequency, however, leads to inferior outcomes. The NIMBUS study (NTR4011) evaluated reduced frequency instillation in a randomized trial of 824 BCG-naïve patients. Patients who received reduced frequency instillation experienced an increased rate of recurrence, at 27% compared with 12% for patients receiving standard BCG treatment (HR 0.40; 95% CI 0.24 to 0.67).<sup>83</sup> While research is ongoing regarding possible alternative dosing regimens, the 6+3 regimen remains the gold standard based on current data.

### BCG-unresponsive NMIBC

BCG-unresponsive NMIBC is a term that encompasses both BCG-refractory and BCG-relapsing (within 6 months of last BCG exposure) NMIBC, as defined by the IBCG.<sup>84</sup> The FDA has also issued guidance on BCG-unresponsive NMIBC with more specific criteria, defined as at least one of the following: (1) persistent or recurrent CIS (with or without recurrent Ta/T1 disease) within 12 months of completion of adequate BCG therapy, (2) recurrent high-grade Ta/T1 disease within 6 months of completion of adequate BCG therapy, or (3) T1 high-grade disease at the first evaluation following BCG induction. Adequate BCG therapy is defined as at least 5 of 6 doses of an initial induction course with at least 2 additional doses (either of maintenance therapy or of a second course of induction).<sup>85</sup> Multiple clinical trials have been conducted using this guidance to define this at-risk patient population, for which the only FDA-approved systemic immunotherapy is pembrolizumab.

Pembrolizumab is approved for the treatment of high-risk, BCG-unresponsive CIS at a dose of 200 mg every 3 weeks or 400 mg every 6 weeks (an alternative treatment schedule approved across multiple indications) for up to 24 months. The approval is based on favorable outcomes for patients in cohort A of the phase II KEYNOTE-057 trial (NCT02625961), who had high-risk, BCG-unresponsive CIS with or without papillary disease and were treated with pembrolizumab every 3 weeks for up to 24 months. Patients were required to undergo full resection of papillary disease prior to their first dose of pembrolizumab. The primary endpoint of CR was assessed at 3 months, and a key secondary endpoint was DOR.<sup>86</sup> At first analysis, 41 of 102 patients achieved CR at 3 months (40.2%; 95% CI 30.6% to 50.4%) and the median DOR of those patients who achieved CR was 12.7 months.<sup>87</sup> At the 12-month landmark analysis for DOR (which occurred approximately 15 months from the start of pembrolizumab therapy), the rate of patients with observed DOR  $\geq 12$  months was 46% (18 of 39 initial complete responders) and 19% of all treated patients.<sup>42</sup> None of the patients developed muscle-invasive or metastatic disease while on protocol treatment. Based on these data, in January 2020, the FDA approved the use of pembrolizumab for the treatment of BCG-unresponsive, high-risk CIS (with or



without papillary tumors).<sup>3</sup> Cohort B of KEYNOTE-057, which focuses on patients with fully resected papillary disease only (without concomitant CIS), was ongoing at the time of publication with a primary endpoint of RFS.<sup>86</sup>

Investigational strategies to overcome BCG-unresponsiveness include concomitant immunomodulation with recombinant cytokines. As an example, N-803 is a mutant IL-15-based immunostimulatory fusion protein complex (IL-15R $\alpha$ Fc) that selectively promotes proliferation and activation of natural killer (NK) cells and CD8<sup>+</sup> T cells. In one cohort of a phase II/III trial that enrolled 80 patients with BCG-unresponsive CIS for intravesical administration of N-803 with BCG, the CR rate at any time was 72% (n=51/71) and the probability of maintaining CR for 12 months was 59%, with a median CR duration of 19.2 months (range 7.6–26.4) months.<sup>88</sup>

A recent phase III, open-label, multicenter US trial (NCT02773849) showed antitumor efficacy with intravesical nadofaragene firadenovec (rAD-IFN $\alpha$ 2b), a replication-defective adenoviral gene transfer vector that delivers interferon- $\alpha$ 2b expression to the bladder epithelium. In the trial, patients with BCG-unresponsive CIS with or without Ta/T1 disease achieved a CR rate at 3 months of 53.4% (95% CI 43.3% to 63.3%) with 45.5% of these remaining free of high-grade recurrence at 12 months. A similar trend of durable response and RFS was observed in patients with papillary high-grade Ta/T1 BCG-unresponsive NMIBC, with 43.8% of patients remaining recurrence-free at 12 months. Progression to muscle invasion occurred in 5% of the CIS cohort and 6% of the high-grade Ta/T1 cohort. Among the patients with CIS, 29% underwent cystectomy by 12 months, as did 21% of those with high-grade Ta/T1 disease. The cystectomy-free survival at 24 months for the whole cohort was 64.5%. Of patients who underwent cystectomy, 3 of 32 (9.3%) in the CIS cohort were found to have pT2 or higher stage disease.<sup>89</sup> While nadofaragene firadenovec is not FDA-approved for the treatment of BCG-unresponsive NMIBC, at the time of writing, it has been granted priority review by the agency and previously received Fast Track and Breakthrough Therapy Designations.

### Panel recommendations

- ▶ BCG is recommended for all eligible patients with high-risk NMIBC (including cases with CIS or papillary tumors) (LE: 1).
- ▶ BCG is also recommended for patients with intermediate-risk NMIBC. However, due to global shortages of BCG, and when BCG is unavailable, the panel recommends intravesical chemotherapy as the first-line therapy for intermediate-risk NMIBC (LE: 1).
- ▶ If patients experience recurrence of intermediate-risk NMIBC after a course of intravesical chemotherapy, the panel recommends BCG as second-line intravesical therapy (LE: 1).
- ▶ BCG is not recommended for the treatment of patients with low-risk NMIBC (LE: 1).

- ▶ BCG should not be administered to patients with active infection or gross hematuria, but BCG may be administered to patients experiencing asymptomatic bacteriuria.
- ▶ Best supportive measures should be employed to ensure that patients receive a full, adequate course of BCG.
- ▶ Pembrolizumab is approved for the treatment of high-risk BCG-unresponsive CIS with or without papillary tumors (LE: 2).

### MUSCLE-INVASIVE BLADDER CANCER

While ICI therapy has not been approved for the treatment of MIBC, a number of ongoing clinical trials are examining the use of ICIs for this disease state. A selection of these trials are summarized in [table 3](#).

#### Phase III trials of neoadjuvant therapy for MIBC

Neoadjuvant cisplatin-based chemotherapy is the current SOC in MIBC. While there are no immunotherapies currently approved as neoadjuvant therapy for localized MIBC, immunotherapy, alone and in combination with chemotherapy, has shown efficacy in MIBC in phase II trials.<sup>40 41 90</sup>

Based on the phase II data, there are several ongoing randomized phase III trials evaluating the role of perioperative immunotherapy in MIBC and results from these trials may establish the utility of this approach. For cisplatin-eligible MIBC, three randomized phase III trials are active at the time of manuscript preparation: KEYNOTE-866 (pembrolizumab, gemcitabine, and cisplatin vs gemcitabine and cisplatin), NIAGARA (durvalumab, gemcitabine, and cisplatin vs gemcitabine and cisplatin), and ENERGIZE (nivolumab, gemcitabine, and cisplatin with or without linrodostat vs gemcitabine and cisplatin). For cisplatin-ineligible MIBC, KEYNOTE-905 is evaluating pembrolizumab with or without EV followed by radical cystectomy versus radical cystectomy alone and the PIVOT IO 009 (NCT04209114) trial is evaluating nivolumab with or without bempegaldesleukin (NKTR-214) followed by radical cystectomy versus radical cystectomy alone. In these trials, immunotherapy agents are continued after radical cystectomy as well. Further, the KEYNOTE-B15 phase III trial, which will examine EV with pembrolizumab versus gemcitabine with cisplatin in patients with cisplatin-eligible MIBC, is anticipated to begin enrollment soon.

#### Phase III trials of adjuvant therapy for MIBC

Atezolizumab was tested in a randomized phase III trial as adjuvant therapy after radical surgery in patients with muscle-invasive (including node-positive) urothelial carcinoma. This trial, IMvigor010, enrolled patients with  $\geq$ ypT2 disease and/or nodal involvement at radical surgery after neoadjuvant chemotherapy or  $\geq$ ypT3 disease and/or nodal involvement if they did not receive prior neoadjuvant chemotherapy. Patients were randomized after radical surgery to receive either atezolizumab for

**Table 3** Ongoing phase III clinical trials of immunotherapy for MIBC

Trial	Immunotherapy and control arms	Agent description	Primary outcome(s) for assessment
<b>Neoadjuvant/adjuvant, cisplatin-eligible</b>			
KEYNOTE-866 (NCT03924856)	Pembrolizumab+gemcitabine+cisplatin Placebo+gemcitabine+cisplatin	ICI, chemotherapy Chemotherapy	pCR rate, EFS
KEYNOTE-B15/EV-304 (NCT04700124)	Pembrolizumab+EV Gemcitabine+cisplatin	ICI, ADC Chemotherapy	pCR rate, EFS
NIAGARA (NCT03732677)	Cisplatin+gemcitabine+durvalumab Cisplatin+gemcitabine	ICI, chemotherapy Chemotherapy	pCR rate at time of surgery, EFS
ENERGIZE CA017078 (NCT03661320)	Gemcitabine+cisplatin Placebo+nivolumab+cisplatin+gemcitabine Linrodostat+nivolumab+cisplatin+gemcitabine	Chemotherapy ICI, chemotherapy IDO1 inhibitor, ICI, chemotherapy	pCR rate, EFS
<b>Neoadjuvant/adjuvant, cisplatin-ineligible</b>			
KEYNOTE-905 (NCT03924895)	Pembrolizumab Surgery alone Pembrolizumab+EV	ICI None ICI, ADC	pCR rate, EFS
PIVOT IO 009 (NCT04209114)	Nivolumab+bempegaldesleukin  Nivolumab Surgery alone	ICI, CD122-biased agonist  ICI None	pCR rate, EFS
<b>Adjuvant</b>			
IMvigor010 (NCT02450331)	Atezolizumab Observation	ICI None	DFS
CheckMate 274 (NCT02632409)	Nivolumab Placebo	ICI None	DFS
AMBASSADOR (NCT03244384)	Pembrolizumab Observation	ICI None	OS, DFS

ADC, antibody-drug conjugate; DFS, disease-free survival; EFS, event-free survival; EV, enfortumab vedotin; ICI, immune checkpoint inhibitor; MIBC, muscle-invasive bladder cancer; OS, overall survival; pCR, pathologic complete response.

1 year or observation. There were no differences in DFS between the arms, which was the primary endpoint. The trial continues to be followed for OS.<sup>44</sup>

Nivolumab has also been evaluated in the adjuvant setting in the randomized placebo-controlled phase III CheckMate 274 trial, which met its co-primary DFS endpoints in the ITT population and the group with tumors with elevated PD-L1 expression in patients who had undergone radical surgery for muscle-invasive (including node-positive) urothelial carcinoma. Initial results from 709 patients (353 randomized to nivolumab including 140 with tumors PD-L1  $\geq 1\%$  and 356 randomized to placebo including 142 with tumors PD-L1  $\geq 1\%$ ) demonstrated a statistically significant and clinically meaningful improvement in median DFS with adjuvant nivolumab after radical surgery compared with placebo, at 21.0 months (range 17.1–33.4) vs 10.9 months (range 8.3–13.9) in the ITT population (HR 0.70; 98.31% CI 0.54 to 0.89;  $p=0.0006$ ) and median DFS not yet reached

(range 22.0 to not estimable) vs 10.8 months (range 5.7 to 21.2) in the tumor PD-L1  $\geq 1\%$  group (HR 0.53; 98.87% CI 0.34 to 0.84;  $p=0.0004$ ). Significant improvement was also seen for the secondary endpoint of non-urothelial tract RFS for adjuvant nivolumab in both the ITT group (HR 0.72; 95% CI 0.58 to 0.89) and in patients with tumors with PD-L  $\geq 1\%$  (HR 0.54; 95% CI 0.38 to 0.77).<sup>45</sup> The FDA granted priority review status to the Biologics License Application for nivolumab for adjuvant treatment of patients with surgically resected, high-risk MIBC in April 2021.

Pembrolizumab is also being tested in a randomized phase III trial as adjuvant therapy after radical surgery in patients with muscle-invasive (node-positive) urothelial carcinoma. This trial, AMBASSADOR, is enrolling patients with  $\geq pT2$  disease and/or nodal involvement at radical surgery after neoadjuvant chemotherapy, or  $\geq pT3$  disease after no neoadjuvant chemotherapy and



**Table 4** Phase II clinical trials of immunotherapy for MIBC

Trial	Interventions	Agent description	Rate of downstaging at time of surgery	Rate of ypT0 at time of surgery
ABACUS (NCT02662309) <sup>41</sup>	Atezolizumab	ICI	NR	31%
PURE-01 (NCT02736266) <sup>151</sup>	Pembrolizumab	ICI	54% (to non-invasive disease)	42%
NABUCCO (NCT03387761) <sup>152</sup>	Nivolumab+ipilimumab	ICI	58% (to non-invasive disease)	45%
NCT02812420 <sup>153</sup>	Durvalumab+tremelimumab	ICI	58% (to ypT1 or less)	38%
BLASST-1 (NCT03294304) <sup>90</sup>	Nivolumab+gemcitabine+cisplatin	ICI+chemotherapy	66%	34%
HCRN GU14-188 (NCT02365766) <sup>154 155</sup>	Pembrolizumab+gemcitabine+cisplatin (eligible cohort) Pembrolizumab+gemcitabine (cisplatin-ineligible cohort)	ICI+chemotherapy	Cisplatin-eligible cohort I: 53% (to ypT0/Tis) Cisplatin-ineligible cohort II: 52% (to ypT0/Tis)	Cisplatin-eligible cohort I: 44% Cisplatin-ineligible cohort II: 45%
DUTRENEO (NCT03472274) <sup>91</sup>	Durvalumab+tremelimumab	ICI	NR	35%

ICI, immune checkpoint inhibitor; MIBC, muscle-invasive bladder cancer; NR, not reported.

will randomize patients to 1 year of pembrolizumab or observation. Accrual is still ongoing.

#### Early phase trials of ICI neoadjuvant therapy for MIBC

Immunotherapy in the neoadjuvant setting has been tested in phase I and II trials, some of which have completed accrual and have reported results (see [table 4](#)). Approaches include ICI as monotherapy before surgery as well as ICI in combination with another ICI or platinum-based chemotherapy. These trials have reported pathologic complete responses (pCRs), also referred to as ypT0, in a subset of patients at the time of surgical intervention, a finding that has correlated with better long-term survival in prior studies of neoadjuvant chemotherapy. Neoadjuvant ICI therapy has been examined in the cisplatin-ineligible (NABUCCO, HCRN GU14-188 Cohort II, NCT02812420) and cisplatin-eligible (BLASST-1, HCRN GU14-188 Cohort I, DUTRENEO) patient populations.

PD-L1 positivity of the tumor by CPS (IHC 22C3 assay) correlated with pembrolizumab responses in the PURE-01 study and ipilimumab and nivolumab in the NABUCCO study. Additionally, in the DUTRENEO trial, for patients in the PD-L1-high group (as measured by the E1L3N XP antibody) the ypT0 rate was 57.1%, while it was 14.3% for patients in the PD-L1-low group.<sup>91</sup> However, all other monotherapy and combination studies in [table 4](#) did not identify a significant response correlation with tumor or IC PD-L1 status.

Overall, ICI monotherapy or combination therapy has been reported to have manageable toxicity and has not been associated with delay of cystectomy. These promising data support ongoing phase III trials, but are not yet practice-changing. Comparisons across these small-moderate sized single-arm phase II trials, however, should be approached cautiously. Significant differences in initial staging techniques, type and length of treatment,

biomarker assays, and eligible patient populations exist across these studies. Additionally, the potential correlation between pathologic assessment after neoadjuvant immunotherapy regimens and OS (and/or RFS) remains unclear.

#### Immunotherapy with chemoradiation as bladder-sparing therapy

Radiation and chemotherapy cause immunogenic cell death,<sup>92 93</sup> which may synergize with ICI therapy to potentiate antitumor responses. Two randomized phase III trials are investigating concurrent anti-PD-(L)1 therapy in combination with external beam radiation and radiosensitizing chemotherapy. The randomized, phase III trial NCT03775265 (SWOG/NRG 1806) is testing atezolizumab and the randomized, phase III KEYNOTE-992 (NCT04241185) is testing pembrolizumab in this setting. In addition, the phase II, randomized NCT03768570 is examining durvalumab following trimodal therapy (surgery, chemotherapy, and radiotherapy).

#### Panel recommendation

- The full results of CheckMate 274 are eagerly awaited to guide the potential use of immunotherapy in the adjuvant setting. Active investigation is ongoing into various neoadjuvant and adjuvant strategies, either as single agents or in combination with chemotherapy, radiotherapy, or novel agents.

#### ADVANCED/METASTATIC UROTHELIAL CARCINOMA

The treatment of mUC typically involves platinum-based chemotherapy as the first-line, SOC modality.<sup>94</sup> Chemotherapy may also play a role in relapsed/refractory (R/R) disease settings. The introduction of immunotherapy, however, has expanded the available options and a number of ICIs have now been approved by the FDA for the treatment of mUC. A treatment algorithm

**Table 5** mUC treatment algorithm

Patient population		Management	
Cisplatin-eligible		Platinum-based chemotherapy	If no disease progression
			Avelumab maintenance
			If disease progression
			Pembrolizumab Avelumab Nivolumab*
Cisplatin-ineligible	PD-L1-positive tumors†	Atezolizumab* Pembrolizumab*	
	PD-L1-negative tumors†	Carboplatin-based chemotherapy	
Cisplatin- and carboplatin-ineligible		Atezolizumab‡* Pembrolizumab‡*	

Individual rows represent treatment decision options that can be followed from left to right horizontally in adjacent columns.

\*Accelerated approvals contingent on confirmatory trials at the time of guideline publication.

†As determined by the appropriate FDA-approved companion diagnostic (ie, PD-L1 staining immune cells (IC)  $\geq 5\%$  of the tumor area by SP142 assay for atezolizumab and combined positive score (CPS)  $\geq 10$  by IHC 22C3 assay for pembrolizumab).

‡Recommendation based on US-only indication.

mUC, advanced/metastatic urothelial cancer; PD-L1, programmed death-ligand 1.

summarizing expert panel recommendations for immunotherapy management of mUC in various patient populations is provided in [table 5](#).

Data from large phase II and III clinical trials evaluating ICIs for mUC are summarized in [table 6](#) and further described in the narrative text below. Another agent, EV, is an ADC that is FDA-approved for the treatment of mUC that has progressed following both platinum-based chemotherapy and ICI treatment.<sup>95</sup>

### Immunotherapies for first-line treatment of mUC

The phase II, single-group assignment clinical trial IMvigor210 examined the efficacy of atezolizumab for the treatment of mUC. In cohort I of the trial (NCT02951767), 119 cisplatin-based chemotherapy-ineligible patients received first-line atezolizumab.<sup>96</sup> The ORR was 24% (95% CI 16% to 32%), and the median DOR was not reached at 2-year follow-up.<sup>31 97</sup> On the basis of ORR and DOR data from both cohorts of the IMvigor210 trial, the FDA approved atezolizumab for the first-line treatment of PD-L1-positive mUC in patients who are ineligible for cisplatin-containing chemotherapy or those who are not eligible for any platinum-based chemotherapy, regardless of PD-L1 status, in April 2017.<sup>5</sup>

In the KEYNOTE-052 (NCT02335424) phase II trial, 374 patients with mUC who were ineligible for cisplatin-based chemotherapy received pembrolizumab as first-line treatment, irrespective of PD-L1 status.<sup>98</sup> At the most recent updated analysis, the ORR was 29% and the median DOR was 30.1 months.<sup>30</sup> Subgroup analysis of the PD-L1-negative (CPS<10) and PD-L1-positive (CPS $\geq$ 10) groups showed higher DOR and OS in the PD-L1-positive group. A follow-up report of long-term outcomes found that 2-year OS was 31.2%.<sup>30</sup> Based on OS and ORR data from KEYNOTE-045 (described in the **Immunotherapies for R/R mUC** section) and ORR and DOR data from KEYNOTE-052, in May 2017, the FDA approved

pembrolizumab for use as a first-line treatment of mUC (in patients who are ineligible for cisplatin-based chemotherapy and PD-L1-positive, or any patient ineligible for platinum-based chemotherapy) and for the treatment of R/R mUC (in patients who have experienced disease progression following platinum-based chemotherapy) regardless of PD-L1 status.<sup>3</sup>

The activity of PD-(L)1 inhibitors for the first-line treatment of cisplatin-ineligible patients with mUC raised the hypothesis of whether PD-(L)1 blockade as single-agent therapy should be extended to cisplatin-eligible patients as well and whether regimens combining platinum-based chemotherapy and PD-(L)1 blockade might further improve outcomes. IMvigor130 was a placebo-controlled phase III trial randomized 1:1:1 to test whether atezolizumab monotherapy or atezolizumab with gemcitabine and platinum (cisplatin or carboplatin) improved survival compared with placebo plus gemcitabine and platinum. The study enrolled 1,213 patients and demonstrated longer PFS for atezolizumab plus chemotherapy compared with chemotherapy alone (8.2 months vs 6.3 months; stratified HR 0.82; 95% CI 0.70 to 0.96; one-sided  $p=0.007$ ). The interim analysis for OS showed a trend toward longer OS with the atezolizumab plus chemotherapy combination, which was not statistically significant (HR 0.83; 95% CI 0.69 to 1.00; one-sided  $p=0.027$ ). The median OS for atezolizumab monotherapy was 15.7 months compared with 13.1 months for chemotherapy. In patients with high levels of PD-L1 expression in tumor-infiltrating ICs (IC2/3 by SP142 assay), atezolizumab monotherapy median OS was not estimable at interim analysis (95% CI 17.7 to not estimable) vs 17.8 months (95% CI 10.0 to not estimable) in the chemotherapy-alone group (stratified HR 0.68; 95% CI 0.43 to 1.08). As atezolizumab versus chemotherapy could not be formally compared at this interim analysis due to the hierarchical

**Table 6** Large phase II and III clinical trials investigating ICIs for mUC

Trial	Design	Results for immunotherapy treatment				
		Interventions (n patients)	ORR	Median DOR	OS	Median PFS
IMvigor210, cohort I (NCT02951767) <sup>31,37</sup>	Phase II, single-arm, open-label	Atezolizumab (first-line) (n=119)	24% (95% CI 16% to 32%)	Median not reached (2-year follow-up)	2-year OS 41% (95% CI 32% to 50%)	2.7 months (95% CI 2.1 to 4.2)
IMvigor210, cohort II (NCT02108652) <sup>17,37</sup>	Phase II, single-arm, open-label	Atezolizumab (R/R) (n=310)	16% (95% CI 13% to 21%)	27.7 months (95% CI 2.1 to 33.4)	2-year OS 23% (95% CI 19% to 28%)	2.1 months (95% CI 2.1 to 2.1)
IMvigor211 (NCT02302807) <sup>28</sup>	Phase III, randomized, open-label	Atezolizumab vs chemotherapy (R/R) (n=931)	PD-L1 IC2/3: 23% ITT: 13.4%	PD-L1 IC2/3: 15.9 months (95% CI 10.4 to NE) ITT: 12.7 months (95% CI 13.0 to 21.7)	Median OS PD-L1 IC2/3: 11.1 months (95% CI 8.6 to 15.5) ITT: 8.6 months (95% CI 7.8 to 9.6) Stratified HR 0.87 (95% CI 0.63 to 1.21; p=0.41)	PD-L1 IC2/3: 2.4 months (95% CI 2.1 to 4.2) ITT: 2.1 months (95% CI 2.1 to 2.2)
IMvigor130 (NCT02807636) <sup>32</sup>	Phase III, randomized, double-blind	Atezolizumab vs chemotherapy (first-line) (n=451)	23% (95% CI 19% to 28%)	NE (95% CI 15.9 months to NE)	Median OS 15.7 months (95% CI 13.1 to 17.8) Stratified HR 0.83 (95% CI 0.69 to 1.00; one-sided p=0.027)	NR
IMvigor130 (NCT02807636) <sup>32</sup>	Phase III, randomized, double-blind	Atezolizumab+gemcitabine/platinum (first-line) vs chemotherapy (n=362)	47% (95% CI 43% to 52%)	8.5 months (95% CI 7.2 to 10.4)	Median OS 16.0 months (95% CI 13.9 to 18.9) HR (1.02 95% CI 0.83 to 1.24)	8.2 months (95% CI 6.5 to 8.3) Stratified HR 0.82 (95% CI 0.70 to 0.96; one-sided p=0.007)
JAVELIN Bladder100 (NCT02603432) <sup>37</sup>	Phase III, randomized, open-label	Avelumab maintenance vs best supportive care (n=700)	9.7% (95% CI 6.8% to 13.3%)	NR	Median OS 21.4 months (95% CI 18.9 to 26.1) HR 0.69 (95% CI 0.56 to 0.86; p=0.001)	3.7 months (95% CI 3.5 to 5.5) HR 0.62 (95% CI 0.52 to 0.75)
DANUBE (NCT02516241) <sup>100</sup>	Phase III, randomized, open-label	Durvalumab+tremelimumab vs chemotherapy (first-line) (n=1,126)	36%	11.1 months (95% CI 7.9 to 18.5)	Median OS 15.1 months (95% CI 13.1 to 18.0) Durvalumab monotherapy HR 0.89 (95% CI 0.71 to 1.11; p=0.30) Durvalumab+tremelimumab HR 0.85 (95% CI 0.72 to 1.02; p=0.075)	NR
CheckMate 275 (NCT02387996) <sup>113</sup>	Phase II, single-arm, open-label	Nivolumab (R/R) (n=386)	20.7% (95% CI 16.1% to 26.1%)	20.3 months (95% CI 11.5 to 31.3)	3-year OS 22%; Median OS 8.6 months (95% CI 6.1 to 11.3)	1.9 months (95% CI 1.9 to 2.3)
KEYNOTE-052 (NCT02335424) <sup>156,157</sup>	Phase II, single-arm, open-label	Pembrolizumab (first-line) (n=374)	24% (95% CI 20% to 29%)	Median not reached (1-year follow-up)	1-year OS 47.5%	2 months (95% CI 2 to 3)
KEYNOTE-045 (NCT02256436) <sup>116</sup>	Phase III, randomized, open-label	Pembrolizumab (R/R) (n=542)	21.1% (95% CI 16.4% to 26.5%)	Median not reached (2-year follow-up)	2-year OS 26.9% HR 0.70 (95% CI 0.57 to 0.85; p<0.001)	2.1 months (95% CI 2.0 to 2.2) HR 0.96 (p=0.32)
KEYNOTE-361 (NCT02853305) <sup>99</sup>	Phase III, randomized, open-label	Pembrolizumab+gemcitabine/platinum vs chemotherapy (first-line) (n=542)	54.7%	8.5 months	1-year OS 61.8% HR 0.86	8.3 months (95% CI 7.5 to 8.5) HR 0.78 (p=0.0033)
HCRN GU14-182 (NCT02500121) <sup>106</sup>	Phase II, randomized, double-blind	Pembrolizumab maintenance vs placebo (n=108)	23%	NR	NR	5.4 months (95% CI 3.1 to 7.3)

CI, confidence interval; DOR, duration of response; HR, hazard ratio; ICI, immune checkpoint inhibitor; ITT, intent-to-treat; mUC, metastatic urothelial cancer; NE, not estimable; NR, not reported; ORR, overall response rate; OS, overall survival; PD-L1, programmed death-ligand 1; PD-L1 IC2/3, PD-L1 expression in ≥5% of tumor-infiltrating immune cells; PFS, progression-free survival; R/R, relapsed/refractory.



statistical analysis plan, these arms will be further evaluated at future analyses. In patients with tumors harboring low PD-L1 expression levels (IC0/1), the median OS was 13.5 months (95% CI 11.1 to 16.4) in the atezolizumab monotherapy group vs 12.9 months (95% CI 11.3 to 15.0) for patients treated with chemotherapy (unstratified HR 1.07; 95% CI 0.86 to 1.33).<sup>32</sup> As noted in the **Diagnostic tests and biomarkers for urothelial cancer immunotherapy** section, however, an unplanned analysis prompted by the Data and Safety Monitoring Committee revealed increased early deaths in patients with low PD-L1 expressing tumors treated with atezolizumab compared with platinum-based chemotherapy—a result that became apparent even before the trial completed accrual and results were publically available.<sup>22 32</sup> This led the FDA and EMA to restrict the label for atezolizumab monotherapy for cisplatin-ineligible patients to only those having tumors with high levels of PD-L1 expression (IC2/3 by Ventana SP142 assay) or, in the US only, patients considered platinum-ineligible (unable to receive even carboplatin) regardless of PD-L1 expression status.

The KEYNOTE-361 trial (NCT02853305) was a phase III randomized trial that assigned 1,010 patients to receive pembrolizumab monotherapy, pembrolizumab with chemotherapy, or SOC chemotherapy. The final results of KEYNOTE-361 revealed no significant improvement in PFS or OS with pembrolizumab plus platinum-based chemotherapy versus platinum-based chemotherapy or with single-agent pembrolizumab versus platinum-based chemotherapy.<sup>33 99</sup> In parallel with events unfolding in the IMvigor130 trial, as described above and in the **Diagnostic tests and biomarkers for urothelial cancer immunotherapy** section, the FDA restricted the prescribing label for pembrolizumab when used in the cisplatin-ineligible frontline setting to patients with PD-L1 expression  $\geq 10$  by CPS or to those who were considered platinum-ineligible (unable to receive carboplatin) regardless of PD-L1 expression. Importantly, neither IMvigor130 nor KEYNOTE-361 were designed to specifically compare single-agent PD-(L)1 blockade versus carboplatin-based chemotherapy in patients ineligible for cisplatin, the current labeled indication for front-line treatment of mUC.

DANUBE (NCT02516241) was a phase III trial that randomized 1,126 patients to receive first-line durvalumab with tremelimumab, durvalumab monotherapy, or SOC chemotherapy. DANUBE did not reach its co-primary endpoints of an improvement in OS for durvalumab versus chemotherapy in patients with PD-L1-high tumors or with tremelimumab plus durvalumab versus chemotherapy in the ITT population.<sup>100</sup>

### Maintenance approaches in MUC

First-line platinum-based chemotherapy results in disease control in mUC in 65%–75% of patients, but PFS and OS are relatively short, with median PFS less than 9 months.<sup>94 101</sup> While significant progress has been made with five anti-PD-(L)1 inhibitors being approved by the

FDA for mUC patients who progress after platinum-based chemotherapy, response rates with second-line anti-PD-(L)1 inhibitors are modest and only a minority of patients obtain durable clinical benefit. Moreover, there is a substantial patient drop off from first-line to second-line therapy in mUC, with only approximately 40% of patients with mUC receiving second-line therapy as seen from patterns of care and outcomes in real-world settings.<sup>102 103</sup>

The current ‘watch-and-wait’ approach for patients with mUC following response or stable disease after first-line platinum-based chemotherapy and prior to initiation of second-line treatment is suboptimal as almost all patients progress within 9 months of active therapy.<sup>94</sup> Maintenance therapy with anti-PD-(L)1 inhibitors after cessation of first-line platinum-based chemotherapy is an attractive treatment strategy in mUC to improve patient outcomes. Historically, maintenance therapy approaches with targeted therapies have not been successful in mUC.<sup>104 105</sup> However, the antitumor activity and relatively favorable safety profile of PD-(L)1 antagonists in mUC make them a potentially attractive option for maintenance therapy.

Two recent trials have evaluated the efficacy of single-agent checkpoint inhibition after completion of first-line systemic platinum-based chemotherapy (HCRN GU14-182 and JAVELIN Bladder 100).<sup>37 106</sup> In the HCRN GU14-182 trial, a phase II study of maintenance pembrolizumab versus placebo after first-line chemotherapy, 108 patients with mUC achieving at least stable disease after up to 8 cycles of first-line platinum-based chemotherapy were enrolled and treated with pembrolizumab or placebo for up to 24 months. The primary endpoint was PFS. Significantly longer PFS was achieved with maintenance pembrolizumab (5.4 months; 95% CI 3.1 to 7.3) compared with placebo (3.0 months; 95% CI 2.7 to 5.5). Median OS was 22 months (95% CI 12.9 to not reached) with pembrolizumab and 18.7 months (95% CI 11.4 to not reached) with placebo, a secondary endpoint for which the trial was not adequately powered and did not reach statistical significance.<sup>106</sup>

The JAVELIN Bladder 100 trial was a phase III, multicenter, multinational, randomized, open-label, parallel-arm study investigating first-line maintenance treatment with avelumab plus best supportive care (BSC) versus BSC alone in patients with mUC who did not have disease progression after first-line platinum-containing chemotherapy. A total of 700 patients were randomly assigned to receive either avelumab plus BSC or BSC alone, and the primary endpoint was OS. Median OS with avelumab and BSC was significantly longer compared with BSC alone (21.4 vs 14.3 months, respectively; HR 0.69; 95% CI 0.56 to 0.86;  $p=0.001$ ). Median PFS was 3.7 months with avelumab and BSC (95% CI 3.5 to 5.5) compared with 2 months with BSC alone (95% CI 1.9 to 2.7; HR 0.56; 95% CI 0.40 to 0.79;  $p<0.001$ ).<sup>38</sup> Based on the results of this trial, the FDA approved avelumab as maintenance therapy for patients with locally advanced or mUC that has not progressed with first-line platinum-containing chemotherapy. It is

important to note that the upfront chemotherapy and maintenance avelumab was approved based a randomized phase III trial, representing a higher LE than the approvals for pembrolizumab and atezolizumab, which were based on phase II data.

### Immunotherapies for R/R mUC

Cohort II of the IMvigor210 trial (NCT02108652) enrolled 310 patients with mUC who had experienced disease progression following platinum-based chemotherapy and explored the activity of atezolizumab.<sup>107</sup> The ORR was 16% (95% CI 13% to 21%), and the median DOR was 27.7 months (95% CI 2.1 to 33.4).<sup>17 97</sup> On the basis of ORR and DOR data from both cohorts of the IMvigor210 trial, the FDA granted accelerated approval to atezolizumab for the treatment of R/R mUC in May 2016.<sup>5</sup>

A randomized open-label phase III trial, IMvigor211, randomized patients with disease progression following platinum-based chemotherapy to either atezolizumab or investigator's choice of chemotherapy (single-agent paclitaxel, docetaxel, or vinflunine (European Union only)). The primary endpoint of this trial was OS in patients with PD-L1-high expression on tumor infiltrating ICs (IC 2/3 by SP142 assay). Of the 931 patients randomized, 234 were PD-L1-high. In that group, there was no significant difference in OS (stratified HR 0.87; 95% CI 0.63 to 1.21;  $p=0.41$ ). Based on the study design, no additional formal analyses were performed, though an OS benefit was observed with atezolizumab versus chemotherapy (regardless of PD-L1 status) in the ITT population in an exploratory analysis. Atezolizumab therapy was associated with fewer grade 3 or 4 AEs and numerically longer DOR than chemotherapy. These results were comparable to those of IMvigor210 cohort II.<sup>28</sup> Because IMvigor211 failed to meet its primary OS endpoint, however, the indication for atezolizumab in patients with mUC who have previously received platinum-based chemotherapy was voluntarily withdrawn in March 2021.

Avelumab received accelerated approval from the FDA in May 2017, for the treatment of R/R mUC based on the JAVELIN Solid Tumor (NCT01772004) phase I expansion trial, which enrolled 242 patients.<sup>4 108</sup> The FDA based its decision on an ORR of 17% (95% CI 11% to 24%) and DOR data (median not reached at 6 months).<sup>26</sup> In an updated safety and efficacy analysis with more than 2 years of follow-up, ORR was 16.5% (95% CI 12.1% to 21.8%), median DOR was 20.5 months (95% CI 9.7 to not reached), and the 24-month OS was 20.1% (95% CI 15.2% to 25.4%) with avelumab.<sup>109</sup>

Study 1108 (NCT01693562), a phase II trial, examined the efficacy of durvalumab in 191 patients with R/R mUC (182 of which had previously received platinum-based chemotherapy).<sup>25 110</sup> The ORR was 20.4% (95% CI 13.1% to 29.5%) and DOR (median not reached at 1 year) data from this trial formed the basis of FDA-accelerated approval for the use of durvalumab to treat

R/R mUC (that had progressed following platinum-based chemotherapy) in May 2017.<sup>111 112</sup> In November 2020, however, the FDA indication for durvalumab for use in previously treated patients with locally advanced or metastatic bladder cancer was voluntarily withdrawn because the phase III DANUBE trial did not meet its primary end points.

Nivolumab was granted accelerated approved by the FDA for the treatment of R/R mUC for patients with disease progression following platinum-based chemotherapy in February 2017.<sup>6</sup> This approval was based on ORR (20.7%; 95% CI 16.1% to 26.1%) and DOR (median 20.3 months; 95% CI 11.5 to 31.3) data from the phase II CheckMate 275 (NCT02387996) trial, which evaluated nivolumab monotherapy in 386 patients with R/R mUC (270 of these patients had experienced disease progression after platinum-based chemotherapy).<sup>113 114</sup> Extended follow-up of this trial confirmed the safety and efficacy data previously reported.<sup>21</sup>

The KEYNOTE-045 (NCT02256436) phase III trial compared pembrolizumab to chemotherapy for 542 patients with mUC who had experienced disease progression after prior platinum-based chemotherapy.<sup>115</sup> At updated long-term follow-up (median 28 months), patients treated with pembrolizumab exhibited a statistically significant advantage compared with chemotherapy in 2-year OS rates (26.9% vs 14.3%; HR 0.70; 95% CI 0.5 to 0.85;  $p=0.00015$ ) and ORR (21.1% vs 11.0%;  $p=0.002$ ). However, there was no significant difference in PFS at the 1-year or 2-year landmarks.<sup>3 116</sup> Median DOR was not reached with pembrolizumab at this updated analysis, but was 4.4 months for chemotherapy.

### Panel recommendations

- The first-line SOC for mUC is platinum-based chemotherapy. Atezolizumab or pembrolizumab can also be considered as first-line therapy for cisplatin-ineligible patients harboring PD-L1-positive tumors based on a companion assay, or for patients who cannot receive carboplatin (the latter in US only) (LE: 2). Combination ICI and chemotherapy treatment are not currently recommended for this setting.
- In patients with with locally advanced or mUC that has not progressed with first-line platinum-containing chemotherapy, avelumab maintenance therapy improves OS (LE: 2).
- Pembrolizumab is recommended for the treatment of patients with platinum-refractory mUC based on a significant OS benefit in a randomized phase III trial (LE: 2). Avelumab and nivolumab also have approvals in this setting.

### IMMUNOTHERAPIES IN DEVELOPMENT FOR UROTHELIAL CANCER

Numerous immunotherapeutic options are currently in advanced stages of development, either alone or

in combination with other agents. Table 7 summarizes information on select novel immunotherapies and immunotherapeutic combinations currently in late phase clinical trials. This table is not intended to be a comprehensive exhaustive list of all trials across therapy settings.

EV is an ADC currently approved as monotherapy for patients with mUC whose cancer has progressed after previous platinum chemotherapy and ICI therapy. EV delivers a payload of monomethyl auristatin E, a tubulin-disrupting agent, to mUC, which overexpresses the nectin-4 surface receptor target of the monoclonal antibody. Although not considered a classical immunotherapy, induction of the innate immune system is emerging as an important mechanism contributing to the antitumor action for EV. Nectin-4 has been found to be a negative regulator of NK cell activity through binding of the inhibitory receptor T cell immunoreceptor with Ig and ITIM domains (TIGIT) and blocking the nectin-4-TIGIT interaction, which enhances NK cell antitumor activity *in vitro* and *in vivo*.<sup>117</sup> Additionally, EV induces an endoplasmic reticulum stress response, which triggers an immunogenic cell death pathway.<sup>118</sup> Several trials are currently studying the potential for enhanced antitumor activity when ADCs are combined with agents that target adaptive antitumor immunity, such as ICIs.<sup>93 119–122</sup> EV103 (cohort A) is a biomarker-agnostic phase Ib trial of first-line EV and pembrolizumab in patients who are cisplatin-ineligible with mUC. Among 43 patients, 93% had a tumor reduction and the ORR was 73%, with a 15% CR rate and a median time to response of 2 months.<sup>122</sup> Responses in this trial were durable; per interim data at median follow-up of 11 months, the median DOR has not been reached.<sup>122</sup> The phase III randomized trial, EV302 (NCT04223856), which randomizes patients to one of three arms of EV and pembrolizumab; EV, pembrolizumab, and platinum-based chemotherapy; or SOC gemcitabine and platinum-based chemotherapy, is currently enrolling.

#### Panel recommendation

- Participation in clinical trials should be discussed with all patients at any stage of bladder cancer.

#### RECOGNITION AND MANAGEMENT OF irAEs

ICIs are associated with a spectrum of irAEs, which may occur in a variety of organ systems, most commonly in the gastrointestinal tract or the skin.<sup>123</sup> While irAEs can generally be managed by temporarily withdrawing ICI treatment and/or with immunosuppressives, such as corticosteroids, severe irAEs may carry significant risks of morbidity or mortality. Given the potential risks associated with ICI treatment, several groups have developed suggested guidelines for the work up and management of suspected irAEs, including the National Comprehensive Cancer Network (NCCN),

American Society of Clinical Oncology (ASCO), European Society of Medical Oncology (ESMO), and SITC.<sup>124–128</sup> While the incidence and severity of irAEs may vary depending on the specific ICI used<sup>129</sup> or on the tumor type being treated, the treatment of urothelial cancer with ICIs has not been shown to carry any risks above baseline for the incidence or severity of irAEs.<sup>130</sup>

#### Panel recommendation

- SITC's guidelines for the management of ICI-related AEs should be consulted for the treatment of irAEs in patients with bladder cancer.

#### PATIENT SUPPORT AND QOL

While immunotherapy for urothelial cancer has well-described benefits for patient outcomes, therapeutic selection and administration also have potential impacts on patient QOL, as well as a unique AE profile. It is important to assess QOL in patients receiving immunotherapy as well as to provide adequate education and support for promptly recognizing and managing any AEs that may occur during or as a result of treatment.

Tools to assess health-related QOL (HRQOL) have been developed that are specific to patients with bladder cancer. Two validated methods of assessment, the Functional Assessment of Cancer Therapy-Vanderbilt Cystectomy Index (FACT-VCI) and the Bladder Cancer Index (BCI), are currently used.<sup>131 132</sup> The FACT-VCI and BCI correlate moderately well, although the BCI is more specifically focused on the effects of bladder cancer treatments.<sup>133</sup> An additional tool to assess outcomes from a patient-focused perspective is the patient-reported outcomes (PRO) version of the Common Terminology Criteria for Adverse Events (PRO-CTCAE). In a study of bladder cancer patients, the PRO-CTCAE showed significant correlation with the European Organization for Research and Treatment of Cancer (EORTC) QLQ-C30, especially for psychological measures.<sup>134</sup>

Despite BCG's long-standing historical efficacy for patients with NMIBC, its administration may negatively affect QOL by causing discomfort and functional impairment. While the majority of patients complete the 6-cycle induction course, a substantial number of patients discontinue maintenance due to symptoms negatively affecting their QOL. In a study of 411 patients with NMIBC treated with BCG, 74.9% were able to complete induction and begin maintenance, but only 52.3% completed 1 year of maintenance treatments. Of the patients who discontinued maintenance, the majority (59.6%) discontinued due to physical discomfort leading to reduced QOL, despite experiencing only grade 1 toxicities.<sup>135</sup> The dose of BCG may also impact QOL—in an analysis of 166 patients, those who received low-dose BCG induction reported significantly improved QOL, with less functional impairment and significantly less occurrence of fever and micturition pain.<sup>136</sup> Nonetheless, further studies are still needed to validate efficacy outcomes of



**Table 7** Immunotherapies in development for the treatment of bladder cancer

Trial	Disease state	Interventions	Agent description	Primary outcome(s) for assessment
S1602 (NCT03091660)	NMIBC (first-line)	TICE-BCG (I/M) Tokyo-172 BCG (I/M) Tokyo-172 BCG (I/M) with priming	BCG strain BCG strain BCG strain	Time to high-grade recurrence
ALBAN (NCT03799835)	NMIBC (first-line)	BCG (I/M) Atezolizumab+BCG (I/M)	BCG ICI, BCG	RFS
POTOMAC (NCT03528694)	NMIBC (high-risk, first-line)	BCG (I/M) Durvalumab+BCG (I/M) Durvalumab+BCG (I)	BCG ICI, BCG ICI, BCG	DFS
B8011006 (NCT04165317)	NMIBC (high-risk, first-line)	BCG (I/M) PF-06801591+BCG (I/M) PF-06801591+BCG (I)	BCG ICI, BCG ICI, BCG	EFS
CheckMate 7G8 (NCT04149574)	NMIBC (high-risk, R/R to BCG)	BCG (I/M)+nivolumab BCG (I/M)+placebo	ICI, BCG BCG	EFS
CheckMate 9UT (NCT03519256)	NMIBC (high-risk, R/R to BCG)	Nivolumab Nivolumab+BCG Nivolumab+BMS-986205 Nivolumab+BMS-986205+BCG	ICI ICI, BCG ICI, IDO1 inhibitor ICI, IDO1 inhibitor, BCG	CR rate, duration of CR
QUILT-3.032 (NCT03022825)	NMIBC (high-risk, R/R to BCG)	BCG+ALT-803 ALT-803	BCG, IL-15 superagonist IL-15 superagonist	CR rate, disease-free rate
MK-3475-676/KEYNOTE-676 (NCT03711032)	NMIBC (high-risk, R/R to BCG)	BCG (I/M)+pembrolizumab BCG (I/M)	ICI, BCG BCG	CR rate
S1605 (NCT02844816)	NMIBC (BCG-unresponsive)	Atezolizumab	ICI	CR rate, EFS
NCT03661320	MIBC (NA, A)	Gemcitabine/cisplatin (NA) Gemcitabine/cisplatin (NA)+nivolumab (NA, A)+placebo (NA, A) Gemcitabine/cisplatin (NA)+nivolumab (NA, A)+BMS-986205 (NA, A)	Chemotherapy ICI, chemotherapy ICI, IDO1 inhibitor, chemotherapy	CR rate, EFS
NIAGARA (NCT03732677)	MIBC (NA, A)	Gemcitabine/cisplatin (NA)+durvalumab (NA, A) Gemcitabine/cisplatin (NA)	ICI, chemotherapy Chemotherapy	CR rate, EFS
MK-3475-905/KEYNOTE-905 (NCT03924895)	MIBC (NA, A)	Pembrolizumab (NA, A) Surgery alone	ICI None	CR rate, EFS
MK-3475-866/KEYNOTE-866 (NCT03924856)	MIBC (NA, A)	Pembrolizumab (NA, A)+gemcitabine/cisplatin (NA) Gemcitabine/cisplatin (NA)+placebo (NA, A)	ICI, chemotherapy Chemotherapy	CR rate, EFS
PIVOT IO 009 (NCT04209114)	MIBC (NA, A)	Nivolumab (NA, A)+NKTR-214 (NA, A) Nivolumab (NA, A) Surgery alone	ICI, CD122-biased agonist ICI None	CR rate, EFS
INTACT SWOG/NRG 1806 (NCT03775265)	MIBC (bladder preservation)	Radiotherapy+chemotherapy Radiotherapy+chemotherapy+atezolizumab	Chemotherapy ICI, chemotherapy	Bladder-intact EFS
MK-3475-992/KEYNOTE-992 (NCT04241185)	MIBC (bladder preservation)	Pembrolizumab+chemotherapy+radiotherapy Placebo+chemotherapy+radiotherapy	ICI, chemotherapy Chemotherapy	Bladder-intact EFS

Continued

Table 7 Continued

Trial	Disease state	Interventions	Agent description	Primary outcome(s) for assessment
AMBASSADOR (NCT03244384)	MIBC (A)	Pembrolizumab Observation	ICI None	OS, DFS
CheckMate 274 (NCT02632409)	MIBC (A)	Nivolumab Placebo	ICI None	DFS
NILE (NCT03682068)	Metastatic (first-line)	Durvalumab+platinum/gemcitabine	ICI, chemotherapy	OS
		Durvalumab+tremelimumab+platinum/gemcitabine	ICI, chemotherapy	
		Platinum/gemcitabine	Chemotherapy	
Checkmate 901 (NCT03036098)	Metastatic (first-line)	Ipilimumab+nivolumab	ICI	OS, PFS
		Ipilimumab+nivolumab+chemotherapy	ICI, chemotherapy	
		Chemotherapy	Chemotherapy	
EV 302 (NCT04223856)	Metastatic (first-line)	EV+pembrolizumab	ICI, ADC	PFS, OS
		Gemcitabine+platinum	Chemotherapy	
		EV+pembrolizumab+platinum	ICI, ADC, chemotherapy	
LEAP-011 (NCT03898180)	Metastatic (first-line)	Pembrolizumab+lenvatinib	ICI, tyrosine kinase inhibitor	PFS, OS
		Pembrolizumab+placebo	ICI	
THOR (NCT03390504)	Metastatic (second- or third-line)	Erdaftinib	FGFR kinase inhibitor	OS
		Pembrolizumab	ICI	

A, adjuvant; ADC, antibody-drug conjugate; AE, adverse event; BCG, Bacillus Calmette-Guérin; CR, complete response; DFS, disease-free survival; EFS, event-free survival; FGFR, fibroblast growth factor receptor; I, induction; ICI, immune checkpoint inhibitor; IL-15, interleukin-15; I/M, induction and maintenance; M, maintenance; MIBC, muscle-invasive bladder cancer; NA, neoadjuvant; NMIBC, non-muscle-invasive bladder cancer; OS, overall survival; PFS, progression-free survival; RFS, recurrence-free survival; R/R, relapsed/refractory.

BCG dose reductions in comparison to full-dose BCG regimens.

While ICIs often exhibit better AE and QOL profiles in comparison to platinum-based chemotherapy, there is a paucity of direct comparisons specifically in patients with bladder cancer. An analysis of participants in the KEYNOTE-045 trial, however, demonstrated that pembrolizumab prolonged time to deterioration in HRQOL score when compared with investigator's choice chemotherapy (median time to deterioration 3.5 months vs 2.3 months, respectively; HR 0.72;  $p=0.004$ ). The change in HRQOL scores from baseline to week 15 of treatment was also significantly different, at 0.69 (95% CI -2.40 to 3.77) in the pembrolizumab arm and -8.36 (95% CI -11.84 to -4.89) in the chemotherapy arm (mean difference 9.05; 95% CI 4.61 to 13.50;  $p<0.001$ ).<sup>137</sup>

Another key consideration for patient QOL is access to treatment and financial limitations. Patients with bladder cancer who report experiencing financial toxicity (defined as 'paying more for medical care than you can afford') may delay treatment due to issues concerning expense management and work productivity. Patients who experience financial toxicity report significantly lower HRQOL.<sup>138</sup> The possibility of financial toxicity presents a major barrier to equitable healthcare access

and contributes to disparities in medical care, including in the context of clinical trial participation.<sup>139 140</sup>

In the context of the SARS-CoV-2 pandemic, cancer care and many aspects of patient QOL may also be impacted. Patients with cancer appear to be more prone to death and to severe outcomes requiring hospitalization from COVID-19.<sup>141-144</sup> Additionally, treatment with ICIs has been associated with increased risk of severe COVID-19 respiratory disease in some analyses, but this association was not confirmed in other analyses.<sup>145-148</sup> The data with regard to BCG treatment is less clear; while some studies indicate that vaccination with BCG (common in regions where tuberculosis infection is a high risk) may be associated with reduced risk of COVID-19,<sup>149 150</sup> further study is still required. Beyond possible impact on outcomes, additional important considerations during the pandemic for patients with cancer include attempts to reduce in-person appointments when possible, use of telemedicine, potential disruptions in public transportation, interruption of clinical trials, and potentially delayed or altered treatment schedules, screening, diagnostic work up, and surveillance. These changes to treatment schedules could create complications in a patient's ability to attend appointments or increase the financial hardship of doing so, potentially negatively impacting cancer care.

## Panel recommendations

- ▶ Patient navigation and PRO tools can help eliminate barriers to oncologic care, enhance patient decision-making, and improve the patient experience during their cancer care. This has been demonstrated in screening outcomes for a variety of malignancies and confirmed in recent studies of NMIBC and MIBC. Combining patient-focused information and educational resources with comprehensive patient-provider conversations can contribute to improved QOL both during treatment and surveillance.
- ▶ Comprehensive conversations with patients about all aspects of medical treatment, including financial obligations, could involve multiple clinical and institutional providers. Conversations should continue throughout patient-provider relationships that reflect the evolving nature of treatment timing, options, and patient concerns.
- ▶ Urothelial cancer-specific outcome measures for BCG and ICI treatments should be developed, validated, and utilized as tools for patient navigation.
- ▶ ICI-specific measures should address a range of treatment protocols and QOL, including ICI alone, combinations with chemotherapy and/or radiation, or any other combination of therapies. Such measures should recognize the often-lengthy nature of bladder cancer treatment and surveillance, along with the potential for adverse effects to occur after the period of initial treatment.
- ▶ Practical patient information and education resources are needed for both BCG and ICI treatment. As more patients are treated with ICIs, written and digital educational materials are needed. Patient information resources in written and digital formats are available from bladder cancer and medical education organizations, in addition to materials provided by the providing clinic.
- ▶ There is now an opportunity to develop, study, and deploy digital/mobile technologies to increase patient awareness and reporting of BCG- and ICI-related AEs. Innovation in patient-provider communication and application of technology to PRO/QOL communication could affect patient care for initial and follow-up of patients with urothelial cancer.

## CONCLUSION

The introduction of ICI therapies has expanded options for patients with urothelial cancer, both in the NMIBC and mUC settings. ICIs, and other immunotherapies, are likely to continue to function as a cornerstone of urothelial cancer treatment, especially as emerging data from ongoing clinical trials provide evidence for their benefits in additional settings. Ongoing clinical trials hold promise for the development of new immunotherapies for the treatment of urothelial cancer, including a gene therapy (nadofaragene firadenovec), a CD122-biased agonist, an IL-15 agonist, an IDO1 inhibitor, and new strains of BCG. The recommendations in this manuscript were based

on available evidence at the time of manuscript preparation and the consensus of the SITC Urothelial Cancer Immunotherapy Guideline Expert Panel. As the field progresses, this guideline will be updated as needed.

## Author affiliations

- <sup>1</sup>Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, New York, New York, USA
- <sup>2</sup>Perlmutter Cancer Center, New York University Langone Medical Center, New York, New York, USA
- <sup>3</sup>Department of Urologic Sciences, The University of British Columbia, Vancouver, British Columbia, Canada
- <sup>4</sup>Department of Genitourinary Medical Oncology, Division of Cancer Medicine, The University of Texas MD Anderson Cancer Center, Houston, Texas, USA
- <sup>5</sup>Bladder Cancer Advocacy Network (BCAN), Bethesda, Maryland, USA
- <sup>6</sup>Dykstra Research, Seattle, Washington, USA
- <sup>7</sup>Department of Medicine, Division of Oncology, University of Washington, Seattle, Washington, USA
- <sup>8</sup>Seattle Cancer Care Alliance, Fred Hutchinson Cancer Research Center, Seattle, Washington, USA
- <sup>9</sup>Department of Hematology and Oncology, Cleveland Clinic Taussig Cancer Institute, Cleveland, Ohio, USA
- <sup>10</sup>Department of Medicine, Duke Cancer Institute, Duke University, Durham, North Carolina, USA
- <sup>11</sup>Division of Hematology-Oncology, Department of Medicine, University of California Los Angeles, Los Angeles, California, USA
- <sup>12</sup>Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA
- <sup>13</sup>The Jesse Brown VA Medical Center, Chicago, Illinois, USA
- <sup>14</sup>Department of Hematology/Oncology, Fox Chase Cancer Center, Philadelphia, Pennsylvania, USA
- <sup>15</sup>Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, New York, USA
- <sup>16</sup>Department of Medicine, Weill Cornell Medical College, New York, New York, USA
- <sup>17</sup>Carolina Urologic Research Center, Myrtle Beach, South Carolina, USA
- <sup>18</sup>Department of Urology and Perlmutter Cancer Center, NYU Langone Medical Center, New York, New York, USA
- <sup>19</sup>Department of Urology under Division of Surgery, The University of Texas MD Anderson Cancer Center, Houston, Texas, USA

**Acknowledgements** The authors acknowledge SITC staff for their contributions including Ben Labbe, PhD and Sam Million-Weaver, PhD for medical writing and Emily Gronseth, PhD; Angela Kilbert; and Lionel Lim for project management and editorial assistance. Additionally, the authors wish to thank the society for supporting the manuscript development. In addition, SITC wishes to thank the Bladder Cancer Advocacy Network (BCAN) for providing a representative to serve on the Urothelial Cancer Immunotherapy Guideline Expert Panel.

**Contributors** All authors served on the SITC Urothelial Cancer Immunotherapy Guideline Expert Panel, drafted content, and provided critical review during the manuscript development. MDG and AMK provided leadership as Chairs of the Expert Panel and provided guidance on the manuscript structure and content and thus are first and last authors; all other authors contributed equally to the manuscript development and review and therefore are listed alphabetically by last name. GSD was the patient advocate representative.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** AVB—Consulting fees: Genentech, Incyte, Janssen, Merck, Pfizer, AstraZeneca, Nektar, Seattle Genetics, DragonFly Therapeutics; Non-CME services: Genentech, AstraZeneca, Merck, Seattle Genetics; Contracted research: Genentech, Merck, AstraZeneca, Nektar, Seattle Genetics, Immunomedics. PCB—Consulting fees: AbbVie, Asieris Pharmaceuticals, Astellas, AstraZeneca, Bayer, Biosynt, Bristol-Myers Squibb, EMD Serono, Ferring, Fergene, H3 Biomedicine, Janssen, Merck, Roche, Sanofi, TerSera, Urogen; Contracted research: iProgen. MTC—Consulting fees: Pfizer, EMD Serono, AstraZeneca, Exelixis, Eisai, Seattle Genetics, Astellas; Contracted research: Exelixis, Pfizer/EMD Serono, Janssen, Apricity Health. GSD—Participant/unpaid volunteer: HealthVibe focus group 2020. MDG—Consulting fees: Inovio, NuMab, Janssen, Merck, GlaxoSmithKline, Genentech, Bristol-Myers Squibb, Seattle Genetics, Dracen, Dragonfly, Astellas, Novartis, Pfizer, AstraZeneca, Incyte, Dendreon, Eli



Lilly, EMD Serono, Aileron Therapeutics; Ownership interest: Rappta Therapeutics; NPI: 1639140668. PG—Consulting fees: AstraZeneca, Dyania Health, EMD Serono, Immunomedics, Janssen, Merck, Mirati Therapeutics, Genentech, Pfizer, Seattle Genetics; Contracted Research: Bavarian Nordic, Bristol-Myers Squibb, Clovis Oncology, Debiopharm, Immunomedics, Pfizer, Merck, QED Therapeutics, GlaxoSmithKline, Kure It Cancer Research, Mirati Therapeutics. SG—Consulting fees: Merck, Bristol-Myers Squibb, Janssen, Seattle Genetics, AstraZeneca; Contracted research: Bristol-Myers Squibb, Pfizer; Non-CME services: Exelixis. CJH—Consulting fees: Astellas, Bristol-Myers Squibb, Genentech, Merck, Eisai, Seattle Genetics, 2bPrecise; Non-CME services: Astellas, Bristol-Myers Squibb, Genentech, Eisai, Seattle Genetics. AMK—Consulting fees: Abbott Molecular, Arquer, ArTara, Asieris, AstraZeneca, BioClin Therapeutics, Bristol-Myers Squibb, Cepheid, Cold Genesys, Eisai, Engene, Inc., Ferring, FerGene, Imagin, Janssen, MDxHealth, Medac, Merck, Pfizer, Photocure, ProTara, Roviant, Seattle Genetics, Sessen Bio, Theralase, TMC Innovation, US Biotest; Contracted research: Adolor, Bristol-Myers Squibb, FDK Industries, Heat Biologics, Merck, Photocure, SWOG/NIH, SPORE, AIBCCR; Clinical trials: FDK, Merck, Bristol-Myers Squibb, Photocure, SWOG, Adolor, Heat Biologics; Laboratory research: NIH, SPORE, AIBCCR; Educational grant: CEC Oncology; Editorial board: European Urology Oncology; Journal of Urology, UroToday; President of organization: International Bladder Cancer Network (IBCN), International Bladder Cancer Group (IBCG); IP Rights: CyPRIT (Cytokine Predictors of Response to Intravesical Therapy) Joint with UT MD Anderson Cancer Center. LPL—Consulting fees: Pfizer Advisory Nursing board, 2018; Lecture/Speaker/Expert Panel: ASCO Bladder Program 2019, Pfizer learning day 2019, ASCO Bladder Program Panel Member and Faculty 2019, Pfizer Learning Day Speaker 2019, BCAN Webinar Faculty 11/2020. JJM—Consulting fees: AstraZeneca, Ferring, Janssen, Nucleix, Foundation Medicine; Contracted research: Epizyme, Tesaro, Abbvie; Teaching an endoscopy course: Olympus. ERP—Consulting fees: AstraZeneca, Infinity Pharma, Pfizer, Flatiron, Genentech, Merck, Seattle Genetics; Contracted research: Astellas, Bristol-Myers Squibb, Genentech, Merck. JER—Consulting fees: Merck, Bristol-Myers Squibb, AstraZeneca, Astellas, Seattle Genetics, Boehringer Ingelheim, Bayer, Pfizer, Roche/Genentech, Mirati, QED Therapeutics, Janssen, Pharmacyclics, GlaxoSmithKline; Contracted research/Clinical trial funding: Seattle Genetics, Astellas, Roche/Genentech, Bayer, AstraZeneca, QED Therapeutics. NS—Consulting fees: Amgen, Astellas, AstraZeneca, Bayer, Bristol-Myers Squibb, Dendreon, Fergene, Ferring, Janssen, Merck, Myovant, Nymox, Pfizer, Sanofi/Genzyme, Sessen Bio, Sun Pharma, Tolmar. GDS—Consulting fees: Merck, Janssen, Ferring, Bristol-Myers Squibb, Pfizer, Genentech, Epivax Oncology, AstraZeneca, Fidia, Cold Genesys, EnGene Bio, Aduro Biotech, Dendreon, FDK Therapies, CiClomed, Seattle Genetics, Urogen, PhotoCure. SITC staff: SMW—Shares owned: Pacific Biosciences, Editas Medicine; EG, AK, BL, LL—Nothing to disclose.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See <http://creativecommons.org/licenses/by-nc/4.0/>.

## REFERENCES

- Siegel RL, Miller KD, Fuchs HE, et al. Cancer statistics, 2021. *CA Cancer J Clin* 2021;71:7–33.
- Woląiewicz M, Hryniewicz R, Grywalska E, et al. Immunotherapy in bladder cancer: current methods and future perspectives. *Cancers* 2020;12:1181.
- Food and Drug Administration, Merck Sharp & Dohme Corp. KEYTRUDA prescribing information. Available: <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm?event=overview.process&AppNo=125514>
- Food and Drug Administration, EMD Serono. BAVENCIO prescribing information. Available: <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm?event=overview.process&AppNo=761049>
- Food and Drug Administration, Genentech Inc. TECENTRIQ prescribing information. Available: <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm?event=overview.process&AppNo=761034>
- Food and Drug Administration, Bristol-Myers Squibb. OPDIVO prescribing information. Available: <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm?event=overview.process&AppNo=125554>
- Challita-Eid PM, Satpayev D, Yang P, et al. Enfortumab Vedotin antibody-drug conjugate targeting Nectin-4 is a highly potent therapeutic agent in multiple preclinical cancer models. *Cancer Res* 2016;76:3003–13.
- Rosenberg J, Sridhar SS, Zhang J, et al. EV-101: a phase I study of single-agent Enfortumab Vedotin in patients with Nectin-4-Positive solid tumors, including metastatic urothelial carcinoma. *J Clin Oncol* 2020;38:1041–9.
- Kamat AM, Bellmunt J, Galsky MD, et al. Society for immunotherapy of cancer consensus statement on immunotherapy for the treatment of bladder carcinoma. *J Immunother Cancer* 2017;5:68.
- Graham R, Mancher M, et al. *Clinical practice guidelines we can trust*. National Academies Press (US), 2011.
- Pietzak EJ, Bagrodia A, Cha EK, et al. Next-Generation sequencing of nonmuscle invasive bladder cancer reveals potential biomarkers and rational therapeutic targets. *Eur Urol* 2017;72:952–9.
- Kamat AM, Briggman J, Urbauer DL, et al. Cytokine panel for response to intravesical therapy (CyPRIT): nomogram of changes in urinary cytokine levels predicts patient response to Bacillus Calmette-Guérin. *Eur Urol* 2016;69:197–200.
- Kamat AM, Karam JA, Grossman HB, et al. Prospective trial to identify optimal bladder cancer surveillance protocol: reducing costs while maximizing sensitivity. *BJU Int* 2011;108:1119–23.
- Lotan Y, Inman BA, Davis LG, et al. Evaluation of the fluorescence in situ hybridization test to predict recurrence and/or progression of disease after Bacillus Calmette-Guérin for primary high grade nonmuscle invasive bladder cancer: results from a prospective multicenter trial. *J Urol* 2019;202:920–6.
- Kamat AM, Willis DL, Dickstein RJ, et al. Novel fluorescence in situ hybridization-based definition of bacille Calmette-Guérin (BCG) failure for use in enhancing recruitment into clinical trials of intravesical therapies. *BJU Int* 2016;117:754–60.
- Haslam A, Prasad V. Estimation of the percentage of US patients with cancer who are eligible for and respond to checkpoint inhibitor immunotherapy drugs. *JAMA Netw Open* 2019;2:e192535.
- Rosenberg JE, Hoffman-Censits J, Powles T, et al. Atezolizumab in patients with locally advanced and metastatic urothelial carcinoma who have progressed following treatment with platinum-based chemotherapy: a single-arm, multicentre, phase 2 trial. *Lancet* 2016;387:1909–20.
- Mariathasan S, Turley SJ, Nickles D, et al. Tgfb attenuates tumour response to PD-L1 blockade by contributing to exclusion of T cells. *Nature* 2018;554:544–8.
- Sharma P, Retz M, Siefker-Radtke A, et al. Nivolumab in metastatic urothelial carcinoma after platinum therapy (CheckMate 275): a multicentre, single-arm, phase 2 trial. *Lancet Oncol* 2017;18:312–22.
- Zhu L, Sun J, Wang L, et al. Prognostic and clinicopathological significance of PD-L1 in patients with bladder cancer: a meta-analysis. *Front Pharmacol* 2019;10:962.
- Galsky MD, Sacci A, Szabo PM, et al. Nivolumab in patients with advanced platinum-resistant urothelial carcinoma: efficacy, safety, and biomarker analyses with extended follow-up from CheckMate 275. *Clin Cancer Res* 2020;26:clincanres.4162.2020.
- Galsky MD, Banchereau R, Hamidi HR, et al. Tumor, immune, and stromal characteristics associated with clinical outcomes with atezolizumab (atezo) + platinum-based chemotherapy (pBC) or atezo monotherapy (mono) versus pBC in metastatic urothelial cancer (mUC) from the phase III IMvigor130 study. *JCO* 2020;38:5011.
- Zajac M, Scott M, Ratcliffe M, et al. Concordance among four commercially available, validated programmed cell death ligand-1 assays in urothelial carcinoma. *Diagn Pathol* 2019;14:99.
- Powles T, Eder JP, Fine GD, et al. MPDL3280A (anti-PD-L1) treatment leads to clinical activity in metastatic bladder cancer. *Nature* 2014;515:558–62.
- Powles T, O'Donnell PH, Massard C, et al. Efficacy and safety of Durvalumab in locally advanced or metastatic urothelial carcinoma: updated results from a phase 1/2 open-label study. *JAMA Oncol* 2017;3:e172411.
- Patel MR, Ellerton J, Infante JR, et al. Avelumab in metastatic urothelial carcinoma after platinum failure (javelin solid tumor): pooled results from two expansion cohorts of an open-label, phase 1 trial. *Lancet Oncol* 2018;19:51–64.
- Herbst RS, Soria J-C, Kowanetz M, et al. Predictive correlates of response to the anti-PD-L1 antibody MPDL3280A in cancer patients. *Nature* 2014;515:563–7.

- 28 Powles T, Durán I, van der Heijden MS, *et al.* Atezolizumab versus chemotherapy in patients with platinum-treated locally advanced or metastatic urothelial carcinoma (IMvigor211): a multicentre, open-label, phase 3 randomised controlled trial. *Lancet* 2018;391:748–57.
- 29 Bellmunt J, de Wit R, Vaughn DJ, *et al.* Pembrolizumab as second-line therapy for advanced urothelial carcinoma. *N Engl J Med* 2017;376:1015–26.
- 30 Vuky J, Balar AV, Castellano D, *et al.* Long-Term outcomes in KEYNOTE-052: phase II study investigating first-line pembrolizumab in Cisplatin-Ineligible patients with locally advanced or metastatic urothelial cancer. *J Clin Oncol* 2020;38:2658–66.
- 31 Balar AV, Galsky MD, Rosenberg JE, *et al.* Atezolizumab as first-line treatment in cisplatin-ineligible patients with locally advanced and metastatic urothelial carcinoma: a single-arm, multicentre, phase 2 trial. *Lancet* 2017;389:67–76.
- 32 Galsky MD, Arija José Ángel Arranz, Bamias A, *et al.* Atezolizumab with or without chemotherapy in metastatic urothelial cancer (IMvigor130): a multicentre, randomised, placebo-controlled phase 3 trial. *Lancet* 2020;395:1547–57.
- 33 Merck. Merck provides update on phase 3 KEYNOTE-361 trial evaluating KEYTRUDA® (pembrolizumab) as monotherapy and in combination with chemotherapy in patients with advanced or metastatic urothelial carcinoma, 2020. Available: <https://www.merck.com/news/merck-provides-update-on-phase-3-keynote-361-trial-evaluating-keytruda-pembrolizumab-as-monotherapy-and-in-combination-with-chemotherapy-in-patients-with-advanced-or-metastatic-urothelial-carc/>
- 34 Galsky MD, ed. *Atezolizumab (atezo) monotherapy versus chemotherapy in previously untreated locally advanced or metastatic urothelial carcinoma (mUC): Clinical outcomes by PD-L1 status in cisplatin (cis)-ineligible pts from the phase III IMvigor130 study*. Genitourinary Cancers Symposium: American Society of Clinical Oncology, 2021.
- 35 Powles T, ed. *1L pembrolizumab (pembro) versus chemotherapy (chemo) for choice-of-carboplatin patients with advanced urothelial carcinoma (UC) in KEYNOTE-361*. Genitourinary Cancers Symposium: American Society of Clinical Oncology, 2021.
- 36 Jiang DM, Gupta S, Kitchlu A, *et al.* Defining cisplatin eligibility in patients with muscle-invasive bladder cancer. *Nat Rev Urol* 2021;18:104–14.
- 37 Powles T, ed. *Maintenance avelumab + best supportive care (BSC) versus BSC alone after platinum-based first-line (1L) chemotherapy in advanced urothelial carcinoma (UC): JAVELIN Bladder 100 phase III interim analysis*. ASCO Virtual Scientific Program: American Society of Clinical Oncology, 2020.
- 38 Powles T, Park SH, Voog E, *et al.* Avelumab maintenance therapy for advanced or metastatic urothelial carcinoma. *N Engl J Med* 2020;383:1218–30.
- 39 Powles T, Park SH, Voog E, *et al.* Maintenance avelumab + best supportive care (BSC) versus BSC alone after platinum-based first-line (1L) chemotherapy in advanced urothelial carcinoma (UC): javelin bladder 100 phase III interim analysis. *JCO* 2020;38:LBA1.
- 40 Necchi A, Raggi D, Gallina A, *et al.* Updated results of PURE-01 with preliminary activity of neoadjuvant pembrolizumab in patients with muscle-invasive bladder carcinoma with variant histologies. *Eur Urol* 2020;77:439–46.
- 41 Powles T, Kockx M, Rodriguez-Vida A, *et al.* Clinical efficacy and biomarker analysis of neoadjuvant atezolizumab in operable urothelial carcinoma in the ABACUS trial. *Nat Med* 2019;25:1706–14.
- 42 Balar AV, Kulkarni GS, Uchio EM, *et al.* Keynote 057: phase II trial of pembrolizumab (pembro) for patients (PTS) with high-risk (HR) nonmuscle invasive bladder cancer (NMIBC) unresponsive to Bacillus Calmette-Guérin (BCG). *JCO* 2019;37:350.
- 43 Black PC, Tangen C, Singh P, *et al.* Phase II trial of atezolizumab in BCG-unresponsive non-muscle invasive bladder cancer: SWOG S1605 (NCT #02844816). *JCO* 2020;38:5022.
- 44 Hussain MHA, Powles T, Albers P, *et al.* IMvigor010: primary analysis from a phase III randomized study of adjuvant atezolizumab (atezo) versus observation (obs) in high-risk muscle-invasive urothelial carcinoma (MIUC). *JCO* 2020;38:5000.
- 45 Bajorin DF, ed. *First results from the phase 3 CheckMate 274 trial of adjuvant nivolumab vs placebo in patients who underwent radical surgery for high-risk muscle-invasive urothelial carcinoma (MIUC)*. Genitourinary Cancers Symposium: American Society of Clinical Oncology, 2021.
- 46 Le DT, Durham JN, Smith KN, *et al.* Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. *Science* 2017;357:409–13.
- 47 Le DT, Uram JN, Wang H, *et al.* Pd-1 blockade in tumors with mismatch-repair deficiency. *N Engl J Med* 2015;372:2509–20.
- 48 Fraune C, Simon R, Hube-Magg C, *et al.* Mmr deficiency in urothelial carcinoma of the bladder presents with temporal and spatial homogeneity throughout the tumor mass. *Urol Oncol* 2020;38:488–95.
- 49 Ju JY, Mills AM, Mahadevan MS, *et al.* Universal Lynch syndrome screening should be performed in all upper tract urothelial carcinomas. *Am J Surg Pathol* 2018;42:1549–55.
- 50 Harper HL, McKenney JK, Heald B, *et al.* Upper tract urothelial carcinomas: frequency of association with mismatch repair protein loss and Lynch syndrome. *Mod Pathol* 2017;30:146–56.
- 51 Metcalfe MJ, Petros FG, Rao P, *et al.* Universal point of care testing for Lynch syndrome in patients with upper tract urothelial carcinoma. *J Urol* 2018;199:60–5.
- 52 Marabelle A, Fakih MG, Lopez J, *et al.* Association of tumour mutational burden with outcomes in patients with select advanced solid tumours treated with pembrolizumab in KEYNOTE-158. *Ann Oncol* 2019;30:v477–8.
- 53 Humphrey PA, Moch H, Cubilla AL, *et al.* The 2016 who classification of tumours of the urinary system and male genital Organs-Part B: prostate and bladder tumours. *Eur Urol* 2016;70:106–19.
- 54 Sylvester RJ, Oosterlinck W, van der Meijden APM. A single immediate postoperative instillation of chemotherapy decreases the risk of recurrence in patients with stage TA T1 bladder cancer: a meta-analysis of published results of randomized clinical trials. *J Urol* 2004;171:2186–90.
- 55 Kamat AM, Witjes JA, Brausi M, *et al.* Defining and treating the spectrum of intermediate risk nonmuscle invasive bladder cancer. *J Urol* 2014;192:305–15.
- 56 Lamm DL, Blumenstein BA, David Crawford E, *et al.* Randomized intergroup comparison of Bacillus Calmette-Guérin immunotherapy and mitomycin C chemotherapy prophylaxis in superficial transitional cell carcinoma of the bladder: a southwest Oncology Group study. *Urol Oncol* 1995;1:119–26.
- 57 Witjes JA, vd Meijden AP, Witjes WP, *et al.* A randomised prospective study comparing intravesical instillations of mitomycin-C, BCG-Tice, and BCG-RIVM in pTa-pT1 tumours and primary carcinoma in situ of the urinary bladder. Dutch South-East cooperative urological group. *Eur J Cancer* 1993;29A:1672–6.
- 58 De Jager R, Guinan P, Lamm D, *et al.* Long-Term complete remission in bladder carcinoma in situ with intravesical TICE Bacillus Calmette Guérin. overview analysis of six phase II clinical trials. *Urology* 1991;38:507–13.
- 59 Food and Drug Administration, Organon Teknika Corp. Tice BCG. Available: <https://www.fda.gov/vaccines-blood-biologics/vaccines/tice-bcg>
- 60 Malmström P-U, Sylvester RJ, Crawford DE, *et al.* An individual patient data meta-analysis of the long-term outcome of randomised studies comparing intravesical mitomycin C versus Bacillus Calmette-Guérin for non-muscle-invasive bladder cancer. *Eur Urol* 2009;56:247–56.
- 61 Sylvester RJ, van der MEIJDEN APM, Lamm DL. Intravesical Bacillus Calmette-Guérin reduces the risk of progression in patients with superficial bladder cancer: a meta-analysis of the published results of randomized clinical trials. *J Urol* 2002;168:1964–70.
- 62 Böhle A, Bock PR. Intravesical Bacille Calmette-Guérin versus mitomycin C in superficial bladder cancer: formal meta-analysis of comparative studies on tumor progression. *Urology* 2004;63:682–6.
- 63 Chou R, Selph S, Buckley DI, *et al.* Intravesical therapy for the treatment of nonmuscle invasive bladder cancer: a systematic review and meta-analysis. *J Urol* 2017;197:1189–99.
- 64 Oddens J, Brausi M, Sylvester R, *et al.* Final results of an EORTC-GU cancers group randomized study of maintenance Bacillus Calmette-Guérin in intermediate- and high-risk TA, T1 papillary carcinoma of the urinary bladder: one-third dose versus full dose and 1 year versus 3 years of maintenance. *Eur Urol* 2013;63:462–72.
- 65 Davies BJ, Hwang TJ, Kesselheim AS. Ensuring Access to Injectable Generic Drugs - The Case of Intravesical BCG for Bladder Cancer. *N Engl J Med* 2017;376:1401–3.
- 66 Messing EM. The BCG shortage. *Bladder Cancer* 2017;3:227–8.
- 67 Bladder Cancer Advocacy Network. Information about the BCG shortage, 2020. Available: <https://bcn.org/2020-information-about-the-bcg-shortage/>
- 68 Chang SS, Boorjian SA, Chou R, *et al.* Diagnosis and treatment of non-muscle invasive bladder cancer: AUA/SUO guideline. *J Urol* 2016;196:1021–9.
- 69 Babjuk M, Böhle A, Burger M, *et al.* EAU guidelines on non-muscle-invasive urothelial carcinoma of the bladder: update 2016. *Eur Urol* 2017;71:447–61.



- 70 Cernuschi T, Malvolti S, Nickels E, *et al.* Bacillus Calmette-Guérin (BCG) vaccine: a global assessment of demand and supply balance. *Vaccine* 2018;36:498–506.
- 71 Babjuk M, Burger M, Compérat EM, *et al.* European Association of Urology Guidelines on Non-muscle-invasive Bladder Cancer (TaT1 and Carcinoma In Situ) - 2019 Update. *Eur Urol* 2019;76:639–57.
- 72 Kamat AM, Porten S. Myths and mysteries surrounding Bacillus Calmette-Guérin therapy for bladder cancer. *Eur Urol* 2014;65:267–9.
- 73 Macleod LC, Ngo TC, Gonzalgo ML. Complications of intravesical Bacillus Calmette-Guérin. *Can Urol Assoc J* 2014;8:540–4.
- 74 Bernini L, Manzini CU, Giuggioli D, *et al.* Reactive arthritis induced by intravesical BCG therapy for bladder cancer: our clinical experience and systematic review of the literature. *Autoimmun Rev* 2013;12:1150–9.
- 75 Durek C, Rüscher-Gerdes S, Jocham D, *et al.* Interference of modern antibacterials with Bacillus Calmette-Guérin viability. *J Urol* 1999;162:1959–62.
- 76 Colombel M, Saint F, Chopin D, *et al.* The effect of ofloxacin on Bacillus Calmette-Guérin induced toxicity in patients with superficial bladder cancer: results of a randomized, prospective, double-blind, placebo controlled, multicenter study. *J Urol* 2006;176:935–9.
- 77 Damiano R, De Sio M, Quarto G, *et al.* Short-Term administration of prulifloxacin in patients with nonmuscle-invasive bladder cancer: an effective option for the prevention of Bacillus Calmette-Guérin-induced toxicity? *BJU Int* 2009;104:633–9.
- 78 Brausi M, Oddens J, Sylvester R, *et al.* Side effects of Bacillus Calmette-Guérin (BCG) in the treatment of intermediate- and high-risk TA, T1 papillary carcinoma of the bladder: results of the EORTC genito-urinary cancers group randomised phase 3 study comparing one-third dose with full dose and 1 year with 3 years of maintenance BCG. *Eur Urol* 2014;65:69–76.
- 79 Bilen CY, Inci K, Erkan I, *et al.* The predictive value of purified protein derivative results on complications and prognosis in patients with bladder cancer treated with Bacillus Calmette-Guérin. *J Urol* 2003;169:1702–5.
- 80 Lüftenegger W, Ackermann DK, Futterlieb A, *et al.* Intravesical versus intravesical plus intradermal Bacillus Calmette-Guérin: a prospective randomized study in patients with recurrent superficial bladder tumors. *J Urol* 1996;155:483–7.
- 81 Saint F, Patard JJ, Irani J, *et al.* Leukocyturia as a predictor of tolerance and efficacy of intravesical BCG maintenance therapy for superficial bladder cancer. *Urology* 2001;57:617–21.
- 82 Martínez-Piñero JA, Flores N, Isorna S, *et al.* Long-Term follow-up of a randomized prospective trial comparing a standard 81 Mg dose of intravesical Bacille Calmette-Guérin with a reduced dose of 27 Mg in superficial bladder cancer. *BJU Int* 2002;89:671–80.
- 83 Grimm M-O, van der Heijden T, Colombel M, *et al.* Treatment of high-grade non-muscle invasive bladder carcinoma by standard number and dose of BCG instillations versus reduced number and standard dose of BCG instillations: results of the phase III clinical trial (NIMBUS). *JCO* 2020;38:436.
- 84 Kamat AM, Sylvester RJ, Böhle A, *et al.* Definitions, end points, and clinical trial designs for non-muscle-invasive bladder cancer: recommendations from the International bladder cancer group. *J Clin Oncol* 2016;34:1935–44.
- 85 Center for Drug Evaluation and Research, Center for Biologics Evaluation and Research. Bacillus Calmette-Guérin-Unresponsive nonmuscle invasive bladder cancer: developing drugs and biologics for treatment guidance for industry, 2018. Available: <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/bacillus-calmette-guerin-unresponsive-nonmuscle-invasive-bladder-cancer-developing-drugs-and>
- 86 Sharp M, Corp D. Study of pembrolizumab (MK-3475) in participants with high risk non-muscle invasive bladder cancer (MK-3475-057/KEYNOTE-057). Available: <https://clinicaltrials.gov/ct2/show/NCT02625961>
- 87 Balar AV. ASCO GU 2019: updated results of Keynote 057: pembrolizumab for patients with high-risk nonmuscle invasive bladder cancer unresponsive to BCG. *ASCO genitourinary cancers symposium; digital: UroToday*, 2019.
- 88 Chamie K, ed. Phase III/III clinical results of IL-15RaFc superagonist N-803 with BCG in BCG-unresponsive non-muscle invasive bladder cancer (NMIBC) carcinoma in situ (CIS) patients. Genitourinary Cancers Symposium: American Society of Clinical Oncology, 2021.
- 89 Boorjian SA, Alemozaffar M, Konety BR, *et al.* Intravesical nadofaragene firadenovec gene therapy for BCG-unresponsive non-muscle-invasive bladder cancer: a single-arm, open-label, repeat-dose clinical trial. *Lancet Oncol* 2021;22:107–17.
- 90 Gupta S, Sonpavde G, Weight CJ, *et al.* Results from BLASST-1 (bladder cancer signal seeking trial) of nivolumab, gemcitabine, and cisplatin in muscle invasive bladder cancer (MIBC) undergoing cystectomy. *JCO* 2020;38:439.
- 91 Grande E, Guerrero F, Puente J, *et al.* DUTRENEO Trial: A randomized phase II trial of DURvalumab and TREmelimumab versus chemotherapy as a NEOadjuvant approach to muscle-invasive urothelial bladder cancer (MIBC) patients (pts) prospectively selected by an interferon (INF)-gamma immune signature. *JCO* 2020;38:5012.
- 92 Galluzzi L, Vitale I, Warren S, *et al.* Consensus guidelines for the definition, detection and interpretation of immunogenic cell death. *J Immunother Cancer* 2020;8:e000337.
- 93 Galluzzi L, Buqué A, Kepp O, *et al.* Immunogenic cell death in cancer and infectious disease. *Nat Rev Immunol* 2017;17:97–111.
- 94 von der Maase H, Sengelov L, Roberts JT, *et al.* Long-Term survival results of a randomized trial comparing gemcitabine plus cisplatin, with methotrexate, vinblastine, doxorubicin, plus cisplatin in patients with bladder cancer. *J Clin Oncol* 2005;23:4602–8.
- 95 Rosenberg JE, O'Donnell PH, Balar AV, *et al.* Pivotal trial of Enfortumab Vedotin in urothelial carcinoma after platinum and Anti-Programmed death 1/Programmed death ligand 1 therapy. *J Clin Oncol* 2019;37:2592–600.
- 96 Roche H-L. A study of Atezolizumab in participants with locally advanced or metastatic urothelial bladder cancer (cohort 1). Available: <https://clinicaltrials.gov/ct2/show/NCT02951767>
- 97 Balar AV, Dreicer R, Loriot Y, *et al.* Atezolizumab (atezo) in first-line cisplatin-ineligible or platinum-treated locally advanced or metastatic urothelial cancer (mUC): long-term efficacy from phase 2 study IMvigor210. *JCO* 2018;36:4523.
- 98 Sharpe M, Corp D. Study of pembrolizumab (MK-3475) in participants with advanced urothelial cancer (MK-3475-052/KEYNOTE-052). Available: <https://clinicaltrials.gov/ct2/show/NCT02335424>
- 99 Alva A, Csöszsi T, Ozguroglu M, *et al.* LBA23 pembrolizumab (P) combined with chemotherapy (C) vs C alone as first-line (1L) therapy for advanced urothelial carcinoma (UC): KEYNOTE-361. *Ann Oncol* 2020;31:S1155–215.
- 100 Powles TB, van der Heijden MS, Castellano Gauna D, *et al.* 697O a phase III, randomized, open-label study of first-line durvalumab (D) with or without tremelimumab (T) vs standard of care chemotherapy in patients with unresectable, locally advanced or metastatic urothelial carcinoma (Danube). *Ann Oncol* 2020;31:S550–1.
- 101 Fisher MD, Shenolikar R, Miller PJ, *et al.* Treatment patterns and outcomes in stage IV bladder cancer in a community oncology setting: 2008-2015. *Clin Genitourin Cancer* 2018;16:e1171–9.
- 102 Galsky MD, Chowdhury S, Bellmunt J, *et al.* Treatment patterns and outcomes in “real world” patients (pts) with metastatic urothelial cancer (UC). *JCO* 2013;31:4525.
- 103 Cheeseman S, Thompson M, Sopwith W, *et al.* Current treatment and outcomes benchmark for locally advanced or metastatic urothelial cancer from a large UK-based single centre. *Front Oncol* 2020;10:167.
- 104 Grivas PD, Daignault S, Tagawa ST, *et al.* Double-Blind, randomized, phase 2 trial of maintenance sunitinib versus placebo after response to chemotherapy in patients with advanced urothelial carcinoma. *Cancer* 2014;120:692–701.
- 105 Powles T, Huddart RA, Elliott T, *et al.* Phase III, double-blind, randomized trial that compared maintenance lapatinib versus placebo after first-line chemotherapy in patients with human epidermal growth factor receptor 1/2-Positive metastatic bladder cancer. *J Clin Oncol* 2017;35:48–55.
- 106 Galsky MD, Mortazavi A, Milowsky MI, *et al.* Randomized double-blind phase II study of maintenance pembrolizumab versus placebo after first-line chemotherapy in patients with metastatic urothelial cancer. *JCO* 2020;38:1797–806.
- 107 Roche H-L. A study of Atezolizumab in participants with locally advanced or metastatic urothelial bladder cancer (cohort 2). Available: <https://clinicaltrials.gov/ct2/show/NCT02108652>
- 108 Sero EMD, KGaA M. Avelumab in Metastatic or Locally Advanced Solid Tumors (JAVELIN Solid Tumor). Available: <https://clinicaltrials.gov/ct2/show/NCT01772004>
- 109 Apolo AB, Ellerton JA, Infante JR, *et al.* Avelumab treatment for metastatic urothelial carcinoma in the phase Ib javelin solid tumor study: updated safety and efficacy analysis with ≥ two years of follow-up. *JCO* 2019;37:425.
- 110 MedImmune. A phase 1/2 study to evaluate MEDI4736. Available: <https://clinicaltrials.gov/ct2/show/NCT01693562>
- 111 Food and Drug Administration, AstraZeneca UK. IMFINZI prescribing information. Available: <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm?event=overview.process&AppNo=761069>



- 112 Powles T, O'Donnell PH, Massard C, *et al.* Updated efficacy and tolerability of durvalumab in locally advanced or metastatic urothelial carcinoma. *JCO* 2017;35:286.
- 113 Siefker-Radtke AO, Baron AD, Necchi A, *et al.* Nivolumab monotherapy in patients with advanced platinum-resistant urothelial carcinoma: efficacy and safety update from CheckMate 275. *JCO* 2019;37:4524.
- 114 Bristol-Myers Squibb, Ono Pharmaceutical. A study of nivolumab in participants with metastatic or unresectable bladder cancer. Available: <https://clinicaltrials.gov/ct2/show/NCT02387996>
- 115 Sharp M, Corp D. A study of pembrolizumab (MK-3475) versus paclitaxel, docetaxel, or vinflunine for participants with advanced urothelial cancer (MK-3475-045/KEYNOTE-045). Available: <https://clinicaltrials.gov/ct2/show/NCT02256436>
- 116 Fradet Y, Bellmunt J, Vaughn DJ, *et al.* Randomized phase III KEYNOTE-045 trial of pembrolizumab versus paclitaxel, docetaxel, or vinflunine in recurrent advanced urothelial cancer: results of >2 years of follow-up. *Ann Oncol* 2019;30:970–6.
- 117 Reches A, Ophir Y, Stein N, *et al.* Nectin4 is a novel TIGIT ligand which combines checkpoint inhibition and tumor specificity. *J Immunother Cancer* 2020;8:e000266.
- 118 Liu BA, Olson D, Snead K. Abstract 5581: Enfortumab vedotin, an anti-Nectin-4 ADC demonstrates bystander cell killing and immunogenic cell death anti-tumor activity mechanisms of action in urothelial cancers. *Cancer Res* 2020;80:5581.
- 119 Gardai SJ, McPhillips KA, Frasca SC, *et al.* Cell-Surface calreticulin initiates clearance of viable or apoptotic cells through trans-activation of LRP on the phagocyte. *Cell* 2005;123:321–34.
- 120 Cao A, Heiser R, Law C-L. Abstract 4914: Auristatin-based antibody drug conjugates activate multiple ER stress response pathways resulting in immunogenic cell death and amplified T-cell responses. *Cancer Res* 2016;76:4914.
- 121 Cao AT, Law C-L, Gardai SJ. Abstract 5588: Brentuximab vedotin-driven immunogenic cell death enhances antitumor immune responses, and is potentiated by PD1 inhibition in vivo. *Cancer Res* 2017;77:5588.
- 122 Hoimes CJ, Rosenberg JE, Srinivas S, *et al.* EV-103: initial results of enfortumab vedotin plus pembrolizumab for locally advanced or metastatic urothelial carcinoma. *Ann Oncol* 2019;30:v356–402.
- 123 Arnaud-Coffin P, Maillet D, Gan HK, *et al.* A systematic review of adverse events in randomized trials assessing immune checkpoint inhibitors. *Int J Cancer* 2019;145:639–48.
- 124 Thompson JA, Schneider BJ, Brahmer J, *et al.* Management of Immunotherapy-Related toxicities, version 1.2019. *J Natl Compr Canc Netw* 2019;17:255–89.
- 125 Brahmer JR, Lacchetti C, Schneider BJ, *et al.* Management of immune-related adverse events in patients treated with immune checkpoint inhibitor therapy: American Society of clinical oncology clinical practice guideline. *J Clin Oncol* 2018;36:1714–68.
- 126 Haanen JBAG, Carbone F, Robert C, Kerr KM, *et al.* Management of toxicities from immunotherapy: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2017;28:iv119–42.
- 127 Puzanov I, Diab A, Abdallah K, *et al.* Managing toxicities associated with immune checkpoint inhibitors: consensus recommendations from the Society for immunotherapy of cancer (SITC) toxicity management Working group. *J Immunother Cancer* 2017;5:95.
- 128 Brahmer JR, Abu-Sbeih H, Ascierto PA, *et al.* Society for immunotherapy of cancer (SITC) clinical practice guideline on immune checkpoint inhibitor-related adverse events. *J Immunother Cancer* 2021;9:e002435.
- 129 Song P, Zhang D, Cui X, *et al.* Meta-Analysis of immune-related adverse events of immune checkpoint inhibitor therapy in cancer patients. *Thorac Cancer* 2020;11:2406–30.
- 130 Wang P-F, Chen Y, Song S-Y, *et al.* Immune-Related adverse events associated with anti-PD-1/PD-L1 treatment for malignancies: a meta-analysis. *Front Pharmacol* 2017;8:730.
- 131 Cookson MS, Dutta SC, Chang SS, *et al.* Health related quality of life in patients treated with radical cystectomy and urinary diversion for urothelial carcinoma of the bladder: development and validation of a new disease specific questionnaire. *J Urol* 2003;170:1926–30.
- 132 Gilbert SM, Dunn RL, Hollenbeck BK, *et al.* Development and validation of the bladder cancer index: a comprehensive, disease specific measure of health related quality of life in patients with localized bladder cancer. *J Urol* 2010;183:1764–70.
- 133 Moncrief TJ, Balaji P, Lindgren BB, *et al.* Comparative evaluation of Bladder-specific health-related quality of life instruments for bladder cancer. *Urology* 2017;108:76–81.
- 134 Taarnhøj GA, Johansen C, Lindberg H, *et al.* Patient reported symptoms associated with quality of life during chemo- or immunotherapy for bladder cancer patients with advanced disease. *Cancer Med* 2020;9:3078–87.
- 135 Serretta V, Scalici Gesolfo C, Alonge V, *et al.* Does the compliance to intravesical BCG differ between common clinical practice and international multicentric trials? *Urol Int* 2016;96:20–4.
- 136 Yokomizo A, Kanimoto Y, Okamura T, *et al.* Randomized controlled study of the efficacy, safety and quality of life with low dose Bacillus Calmette-Guérin instillation therapy for nonmuscle invasive bladder cancer. *J Urol* 2016;195:41–6.
- 137 Vaughn DJ, Bellmunt J, Fradet Y, *et al.* Health-Related quality-of-life analysis from KEYNOTE-045: a phase III study of pembrolizumab versus chemotherapy for previously treated advanced urothelial cancer. *J Clin Oncol* 2018;36:1579–87.
- 138 Casilla-Lennon MM, Choi SK, Deal AM, *et al.* Financial toxicity among patients with bladder cancer: reasons for delay in care and effect on quality of life. *J Urol* 2018;199:1166–73.
- 139 Chino F, Zafar SY. Financial toxicity and equitable access to clinical trials. *Am Soc Clin Oncol Educ Book* 2019;39:11–18.
- 140 Nipp RD, Lee H, Gorton E, *et al.* Addressing the financial burden of cancer clinical trial participation: longitudinal effects of an equity intervention. *Oncologist* 2019;24:1048–55.
- 141 Yu J, Ouyang W, Chua MLK, *et al.* SARS-CoV-2 transmission in patients with cancer at a tertiary care hospital in Wuhan, China. *JAMA Oncol* 2020;6:1108–10.
- 142 Liang W, Guan W, Chen R, *et al.* Cancer patients in SARS-CoV-2 infection: a nationwide analysis in China. *Lancet Oncol* 2020;21:335–7.
- 143 Dai M, Liu D, Liu M, *et al.* Patients with cancer appear more vulnerable to SARS-CoV-2: a multicenter study during the COVID-19 outbreak. *Cancer Discov* 2020;10:783–91.
- 144 Hijano DR, Maron G, Hayden RT. Respiratory viral infections in patients with cancer or undergoing hematopoietic cell transplant. *Front Microbiol* 2018;9:3097.
- 145 Robilotti EV, Babady NE, Mead PA, *et al.* Determinants of COVID-19 disease severity in patients with cancer. *Nat Med* 2020;26:1218–23.
- 146 Kuderer NM, Choueiri TK, Shah DP, *et al.* Clinical impact of COVID-19 on patients with cancer (CCC19): a cohort study. *Lancet* 2020;395:1907–18.
- 147 Grivas P, Warner J, Shyr Y, *et al.* LBA72 assessment of clinical and laboratory prognostic factors in patients with cancer and SARS-CoV-2 infection: the COVID-19 and cancer Consortium (CCC19). *Ann Oncol* 2020;31:S1202–3.
- 148 Wise-Draper TM, Desai A, Elkrief A, *et al.* LBA71 systemic cancer treatment-related outcomes in patients with SARS-CoV-2 infection: a CCC19 registry analysis. *Ann Oncol* 2020;31:S1201–2.
- 149 Redelman-Sidi G. Could BCG be used to protect against COVID-19? *Nat Rev Urol* 2020;17:316–7.
- 150 Hegarty PK, Sfakianos JP, Giannarini G, *et al.* COVID-19 and Bacillus Calmette-Guérin: what is the link? *Eur Urol Oncol* 2020;3:259–61.
- 151 Necchi A, Anichini A, Raggi D, *et al.* Pembrolizumab as neoadjuvant therapy before radical cystectomy in patients with muscle-invasive urothelial bladder carcinoma (PURE-01): an open-label, single-arm, phase II study. *J Clin Oncol* 2018;36:3353–60.
- 152 Van der Heijden MS, van Dijk N, Smit L, *et al.* Pre-Operative ipilimumab and nivolumab in locoregionally advanced, stage III, urothelial cancer (NABUCCO). *Ann Oncol* 2019;30:v358–402.
- 153 Gao J, Navai N, Alhalabi O, *et al.* Neoadjuvant PD-L1 plus CTLA-4 blockade in patients with cisplatin-ineligible operable high-risk urothelial carcinoma. *Nat Med* 2020;26:1845–51.
- 154 Hoimes CJ, Adra N, Fleming MT, *et al.* Phase Ib/II neoadjuvant (N-) pembrolizumab (P) and chemotherapy for locally advanced urothelial cancer (laUC): Final results from the cisplatin (C)- eligible cohort of HCRN GU14-188. *JCO* 2020;38:5047.
- 155 Kaimakliotis HZ, Adra N, Kelly WK, *et al.* Phase II neoadjuvant (N-) gemcitabine (G) and pembrolizumab (P) for locally advanced urothelial cancer (laUC): Interim results from the cisplatin (C)- ineligible cohort of GU14-188. *JCO* 2020;38:5019.
- 156 Balar AV, Castellano D, O'Donnell PH, *et al.* First-Line pembrolizumab in cisplatin-ineligible patients with locally advanced and unresectable or metastatic urothelial cancer (KEYNOTE-052): a multicentre, single-arm, phase 2 study. *Lancet Oncol* 2017;18:1483–92.
- 157 Vuky J, Balar AV, Castellano DE, *et al.* Updated efficacy and safety of KEYNOTE-052: a single-arm phase 2 study investigating first-line pembrolizumab (pembro) in cisplatin-ineligible advanced urothelial cancer (UC). *JCO* 2018;36:4524.